



Rabbit Anti-Phospho-Ack1 (Tyr326) antibody

SL3046R

Product Name:	Phospho-Ack1 (Tyr326)
Chinese Name:	磷酸化Ack1抗体
Alias:	Acetate kinase 1; Acetokinase 1; Activated CDC42 kinase 1; Activated Cdc42 associated kinase 1; Activated p21cdc42Hs kinase; Tyrosine kinase non receptor 2; Tyrosine kinase non-receptor protein 2; non-receptor protein tyrosine kinase ACK; ACK; TNK 2; Tnk2; ACK1; FLJ44758; FLJ45547; ACK 1; ACK-1; ACK1_MOUSE; Activated p21cdc42Hs kinase; FLJ44758; FLJ45547; Tyrosine kinase non receptor protein 2; p21cdc42Hs.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Dog,Pig,Cow,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	117kDa
Cellular localization:	The nucleuscytoplasmicThe cell membrane
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated Synthesised phosphopeptide derived from human Ack1 around the phosphorylation site of Tyr326:FT(p-Y)GQ
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed

This gene encodes a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The protein may be involved in a regulatory mechanism that sustains the GTP-bound active form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation signal transduction pathway. Several alternatively spliced transcript variants have been identified from this gene, but the full-length nature of only two transcript variants has been determined. [provided by RefSeq.]

Function:

Non-receptor tyrosine-protein and serine/threonine-protein kinase that is implicated in cell spreading and migration, cell survival, cell growth and proliferation. Transduces extracellular signals to cytosolic and nuclear effectors. Phosphorylates AKT1, AR, MCF2, WASL and WWOX. Implicated in trafficking and clathrin-mediated endocytosis through binding to epidermal growth factor receptor (EGFR) and clathrin. Binds to both poly- and mono-ubiquitin and regulates ligand-induced degradation of EGFR, thereby contributing to the accumulation of EGFR at the limiting membrane of early endosomes. Downstream effector of CDC42 which mediates CDC42-dependent cell migration via phosphorylation of BCAR1. May be involved both in adult synaptic function and plasticity and in brain development. Activates AKT1 by phosphorylating it on 'Tyr-176'. Phosphorylates AR on 'Tyr-267' and 'Tyr-363' thereby promoting its recruitment to androgen-responsive enhancers (AREs). Phosphorylates WWOX on 'Tyr-287'. Phosphorylates MCF2, thereby enhancing its activity as a guanine nucleotide exchange factor (GEF) toward Rho family proteins. Contributes to the control of AXL receptor levels. Confers metastatic properties on cancer cells and promotes tumor growth by negatively regulating tumor suppressor such as WWOX and positively regulating pro-survival factors such as AKT1 and AR.

Subunit:

Homodimer. Interacts with CSPG4 (activated). Interacts with MERTK (activated); stimulates autophosphorylation. May interact (phosphorylated) with HSP90AB1; maintains kinase activity. Interacts with NPHP1. Interacts with SNX9 (via SH3 domain). Interacts with SRC (via SH2 and SH3 domain). Part of a collagen stimulated complex involved in cell migration composed of CDC42, CRK, TNK2 and BCAR1/p130cas. Interacts with BCAR1/p130cas via SH3 domains. Forms complexes with GRB2 and numerous receptor tyrosine kinases (RTK) including LTK, AXL or PDGFRL, in which GRB2 promotes RTK recruitment by TNK2 (By similarity). Interacts with CDC42. Interacts with EGFR, and this interaction is dependent on EGF stimulation and kinase activity of EGFR. Interacts (via kinase domain) with AKT1. Interacts with NEDD4 (via WW3 domain). NEDD4L and EGF promote association with NEDD4.

Subcellular Location:

Cell membrane. Nucleus. Endosome. Cell junction, adherens junction. Cytoplasmic vesicle membrane; Peripheral membrane protein; Cytoplasmic side (By similarity). Cytoplasmic vesicle, clathrin-coated vesicle (By similarity). Membrane, clathrin-coated

Product Detail:

pit (By similarity). Note=The Tyr-284 phosphorylated form is found both in the membrane and nucleus. Co-localizes with EGFR on endosomes. Nuclear translocation is CDC42-dependent.

Tissue Specificity:

Ubiquitously present in all tissues tested. Highly expressed in the adult central nervous system (CNS); hippocampus, neocortex, and cerebellum, both at dendritic spines and presynaptic axon terminals. Levels are strongly increased during enhanced neural activity.

Post-translational modifications:

Autophosphorylation regulates kinase activity. Phosphorylation on Tyr-533 is required for interaction with SRC and is observed during association with clathrin-coated pits (By similarity).

Polyubiquitinated by NEDD4 and NEDD4L. Degradation can be induced by EGF and is lysosome-dependent.

Similarity:

Belongs to the protein kinase superfamily. Tyr protein kinase family.

Contains 1 CRIB domain.

Contains 1 protein kinase domain.

Contains 1 SH3 domain.

Contains 1 UBA domain.

SWISS:

Q07912

Gene ID:

10188

Database links:

[Entrez Gene: 10188](#)Human

[Entrez Gene: 51789](#)Mouse

[Entrez Gene: 280710](#)Cow

[Entrez Gene: 303882](#)Rat

[Omim: 606994](#)Human

[SwissProt: Q17R13](#)Cow

[SwissProt: Q07912](#)Human

[SwissProt: O54967](#)Mouse

[SwissProt: Q5U2X5](#)Rat

[Unigene: 518513](#)Human

[Unigene: 251115](#)Mouse

[Unigene: 98335](#)Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Ack

1分子是一组促进酪氨酸激酶生长的家族成员, 该蛋白与Tumour分裂、生长、转移起到很重要的作用, 可加速恶性Tumour的浸润。

www.sunlongbiotech.com