

Rabbit Anti-phospho-Afadin (Ser1721) antibody

SL3072R

| Product Name: | phospho-Afadin (Ser1721) |
|------------------------|--|
| Chinese Name: | 磷酸化丝状肌动蛋白Binding protein抗体 |
| Alias: | p-Afadin(Ser1721); Afadin (phospho S1721); Afadin (phospho Ser1721); MLLT4; Protein AF6; AF6; Afadin; AFAD_HUMAN; ALL1-fused gene from chromosome 6 protein; Protein AF-6. |
| | D 113 |
| Organism Species: | Rabbit |
| Clonality: | Polyclonal |
| React Species: | Human, Mouse, Rat, |
| Applications: | WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. |
| Molecular weight: | 207kDa |
| Cellular localization: | The cell membrane |
| Form: | Lyophilized or Liquid |
| Concentration: | lmg/ml |
| immunogen: | KLH conjugated Synthesised phosphopeptide derived from human Afadin around the phosphorylation site of Ser1721:VL(p-S)PD- |
| Lsotype: | IgG |
| Purification: | affinity purified by Protein A |
| Storage Buffer: | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. |
| Storage: | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |
| PubMed: | PubMed |
| Product Detail: | Afadin is a cell-cell adherens junction F-actin binding multidomain protein. Two splice variants of afadin have been described: l-afadin, which is ubiquitously expressed, and the smaller s-afadin, which is abundantly expressed in neural tissue. l-Afadin contains |

one PDZ domain near its middle followed by three proline-rich and one F-actin-binding region at the carboxyl-terminal. l/s-Afadin is localized to cell-cell junctions and binds several cytoplasmic proteins such as the small GTPase Ras, the tight junction protein ZO-1, and the vinculin binding protein ponsin/SH3P12.

Function:

Belongs to an adhesion system, probably together with the E-cadherin-catenin system, which plays a role in the organization of homotypic, interneuronal and heterotypic cell-cell adherens junctions (AJs). Nectin- and actin-filament-binding protein that connects nectin to the actin cytoskeleton.

Subunit:

Homodimer. Interacts with F-actin, nectin and PVRL3/nectin-3. Essential for the association of nectin and E-cadherin. Isoform 1/s-afadin does not interact with F-actin. Interacts with ZO-1 and occludin, but probably in an indirect manner. Interacts with RIT1 and RIT2 (By similarity). Interacts with NRXN1 and BCR.

Subcellular Location:

Cell junction, adherens junction. Note=Not found at cell-matrix AJs.

DISEASE:

Note=A chromosomal aberration involving MLLT4 is associated with acute leukemias. Translocation t(6;11)(q27;q23) with MLL/HRX. The result is a rogue activator protein.

Similarity:

Contains 1 dilute domain.

Contains 1 FHA domain.

Contains 1 PDZ (DHR) domain.

Contains 2 Ras-associating domains

SWISS:

P55196

Gene ID:

4301

Database links:

Entrez Gene: 4301Human

Entrez Gene: 17356 Mouse

Entrez Gene: 26955Rat

Omim: 159559Human

SwissProt: P55196Human

SwissProt: Q9QZQ1Mouse

SwissProt: O35889Rat

Unigene: 614974Human

Unigene: 644024Human

Unigene: 708726Human

Unigene: 728849Human

Unigene: 59167Mouse

Unigene: 58Rat

Important Note:

MNN SURIORODIOTE

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.