



Rabbit Anti-Phospho-PEA15 (Ser104) antibody

SL3328R

Product Name:	Phospho-PEA15 (Ser104)
Chinese Name:	磷酸化星形胶质细胞PEA15抗体
Alias:	PEA15(Phospho Ser104); PEA15 (Phospho-S104); p-PEA15 (Ser104); Astrocytic phosphoprotein PEA 15; Astrocytic phosphoprotein PEA15; HMAT 1; HMAT1; Homolog of mouse MAT 1 oncogene; Homolog of mouse MAT1 oncogene; HUMMAT 1H; HUMMAT1H; MAT 1; MAT 1H; MAT1; MAT1H; PEA 15; PEA-15; PEA15 protein; PED; Phosphoprotein enriched in astrocytes 15; Phosphoprotein enriched in astrocytes 15kD; Phosphoprotein enriched in diabetes.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Dog,Cow,Rabbit,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	15kDa
Cellular localization:	cytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated Synthesised phosphopeptide derived from human PEA15 around the phosphorylation site of Ser104:IP(p-S)AK
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed:	PubMed
Product Detail:	<p>PED/PEA 15 (Phosphoprotein Enriched in Diabetes/Phosphoprotein Enriched in Astrocytes 15 kDa) is a widely expressed 15 kDa protein comprised of an N terminal region containing a canonical Death Effector Domain (DED) sequence and a nuclear export signal, and a C terminal region containing two serine phosphorylation sites. PED/PEA 15 has been implicated in the regulation of multiple cellular processes including apoptosis, integrin activation, and insulin sensitive glucose transport in insulin responsive cells. Phosphorylation of both serine 104 (a Protein Kinase C site) and serine 116 (a substrate of CaMKII and Akt) is required for PED/PEA 15 function.</p> <p>Subunit: Binds RPS6KA3, MAPK3 and MAPK1. Transient interaction with PLD1 and PLD2 (By similarity). Interacts with CASP8 and FADD.</p> <p>Subcellular Location: Cytoplasm.</p> <p>Tissue Specificity: Ubiquitously expressed. Most abundant in tissues such as heart, brain, muscle and adipose tissue which utilize glucose as an energy source. Lower expression in glucose-producing tissues. Higher levels of expression are found in tissues from individuals with type 2 diabetes than in controls.</p> <p>Similarity: Contains 1 DED (death effector) domain.</p> <p>SWISS: Q15121</p> <p>Gene ID: 8682</p> <p>Database links: Entrez Gene: 8682Human Entrez Gene: 18611Mouse Omim: 603434Human SwissProt: Q15121Human SwissProt: Q62048Mouse Unigene: 517216Human Unigene: 544Mouse</p>

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

www.sunlongbiotech.com