



Rabbit Anti-Phospho-PKR (Thr446 + Thr451) antibody

SL3337R

Product Name:	Phospho-PKR (Thr446 + Thr451)
Chinese Name:	磷酸化蛋白激酶R抗体
Alias:	double-stranded RNA-dependent Protein Kinase; interferon-induced, double-stranded RNA-activated protein kinase isoform a; protein kinase, interferon-inducible double stranded RNA dependent; interferon-inducible eIF2alpha kinase; double stranded RNA activated protein kinase; p68 kinase; eIF-2A protein kinase 2; P1/eIF-2A protein kinase; protein kinase RNA-activated; interferon-inducible RNA-dependent protein kinase; EIF2AK2; EIF2AK1; MGC126524; PKR; PRKR; E2AK2 HUMAN.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	62kDa
Cellular localization:	The nucleuscytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthesised phosphopeptide derived from human PKR around the phosphorylation site of Thr446/451:KR(p-T)RSKG(p-T)LR
Isotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of

	antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	<p>PKR is an interferon-inducible serine/threonine specific protein kinase. It is widely expressed in eukaryotic organisms and activated by double stranded RNA. Activation of PKR by dsRNAs leads to autophosphorylation at multiple sites. Phosphorylation of Thr446 and Thr451 in the PKR activation loop is required in vivo and in vitro for high level kinase activity. PKR phosphorylates its natural substrate, the alpha subunit of eukaryotic protein synthesis initiation factor 2 (EIF2 alpha), leading to the inhibition of protein synthesis. PKR is also involved in TLR signaling and mediates apoptosis in fibroblasts in response to viral infection and inflammatory cytokines, and also activates IKK and NFkB, thereby suppressing apoptosis. Recently, it has been reported that PKR also phosphorylates human p53 on serine 392. PKR might play a role in ER stress-induced apoptosis and in Alzheimer's disease. Alzheimer cases show prominent PKR activation in association with neuritic plaques and pyramidal neurons in the hippocampus and neocortex.</p> <p>Function: Following activation by double-stranded RNA in the presence of ATP, the kinase becomes autophosphorylated and can catalyze the phosphorylation of the translation initiation factor EIF2S1, which leads to an inhibition of the initiation of protein synthesis. Double-stranded RNA is generated during the course of a viral infection. In addition to serine/threonine-protein kinase activity, also has tyrosine-protein kinase activity: phosphorylates CDK1 upon DNA damage. CDK1 phosphorylation triggers CDK1 polyubiquitination and subsequent proteolysis, thus leading to G2 arrest</p> <p>Subunit: Homodimer. Interacts with STRBP (By similarity). Interacts with DNAJC3. Inhibited by direct interaction with viral proteins such as HCV E2, HCV NS5A and influenza A NS1. Activated by the interaction with HIV-1 Tat (By similarity). Forms a complex with FANCA, FANCC, FANCG and HSP70.</p> <p>Post-translational modifications: Autophosphorylated on several Ser and Thr residues. Autophosphorylation of Thr-451 is dependent on Thr-446 and is stimulated by dsRNA binding and dimerization. Autophosphorylation apparently leads to the activation of the kinase.</p> <p>Similarity: Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. GCN2 subfamily. Contains 2 DRBM (double-stranded RNA-binding) domains. Contains 1 protein kinase domain.</p> <p>SWISS: P19525</p> <p>Gene ID:</p>

5610

Database links:

[Entrez Gene: 5610](#)Human

[Omin: 176871](#)Human

[SwissProt: P19525](#)Human


[SwissProt: Q52M43](#)Human

[Unigene: 131431](#)Human

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Picture:



Sample:


Liver (Mouse) Lysate at 40 ug

Primary: Anti- p-PKR (Thr446 + Thr451) (SL3337R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 62 kD

Observed band size: 62 kD



Sample:


Spleen (Mouse) Lysate at 40 ug

Primary: Anti- p-PKR (Thr446 + Thr451) (SL3337R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 62 kD

Observed band size: 62 kD



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-Phospho-PKR(Thr446+Thr451) Polyclonal Antibody, Unconjugated(SL3337R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining