



Rabbit Anti-Phospho-RPS6 (Ser235+Ser236) antibody

SL3388R

| | |
|-------------------------------|---|
| Product Name: | Phospho-RPS6 (Ser235+Ser236) |
| Chinese Name: | 磷酸化S6核糖体蛋白抗体 |
| Alias: | RPS6 (phospho S235 + S236); p-RPS6 (phospho S235 + S236); Air8; DS6; Hemocytes enlarged; Hen; L(1)AIR8; Lethal (1) haemocytes enlarged; Lethal aberrant immune response 8; Lethal(1)abberant immune response[8]; Phosphoprotein NP33; Pp30; Ribosomal protein S6; RP S6; RPS 6; RpS6; RS6; RS6_HUMAN; S6; S6 Ribosomal Protein. |
| Organism Species: | Rabbit |
| Clonality: | Polyclonal |
| React Species: | Human,Mouse,Rat,Chicken,Dog,Pig,Rabbit, |
| Applications: | ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. |
| Molecular weight: | 27kDa |
| Cellular localization: | The nucleuscytoplasmic |
| Form: | Lyophilized or Liquid |
| Concentration: | 1mg/ml |
| immunogen: | KLH conjugated Synthesised phosphopeptide derived from human S6 Ribosomal Protein around the phosphorylation site of Ser235/236:RL(p-S)(p-S)LR |
| Lsotype: | IgG |
| Purification: | affinity purified by Protein A |
| Storage Buffer: | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. |
| Storage: | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |

PubMed:[PubMed](#)

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein kinases. Phosphorylation is induced by a wide range of stimuli, including growth factors, tumor-promoting agents, and mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008]

Function:

May play an important role in controlling cell growth and proliferation through the selective translation of particular classes of mRNA.

Post-translational modifications:

Ribosomal protein S6 is the major substrate of protein kinases in eukaryote ribosomes. The phosphorylation is stimulated by growth factors, tumor promoting agents, and mitogens. It is dephosphorylated at growth arrest. Phosphorylated at Ser-235 and Ser-236 by RPS6KA1 and RPS6KA3; phosphorylation at these sites facilitates the assembly of the preinitiation complex.

Product Detail:**Similarity:**

Belongs to the ribosomal protein S6e family.

SWISS:

P62753

Gene ID:

6194

Database links:

[Entrez Gene: 6194](#) Human

[Entrez Gene: 20104](#) Mouse

[Entrez Gene: 667739](#) Mouse

[Entrez Gene: 29304](#) Rat

[Omim: 180460](#) Human

[SwissProt: P62753](#) Human

[SwissProt: P62754](#) Mouse

[SwissProt: P62755](#) Rat

[Unigene: 408073](#) Human

[Unigene: 379007](#) Mouse


[Unigene: 425090](#) Mouse

[Unigene: 34341](#) Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Picture:



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-RPS6 (Ser235+Ser236)) Polyclonal Antibody, Unconjugated (SL3388R) at 1:500 overnight at 4°C, followed by a

| | |
|--|---|
| | conjugated secondary (sp-0023) for 20 minutes and DAB staining. |
|--|---|

www.sunlongbiotech.com