



Rabbit Anti-Phospho-SRC3 (Thr24) antibody

SL3408R

Product Name:	Phospho-SRC3 (Thr24)
Chinese Name:	磷酸化类固醇受体辅助活化因子-3
Alias:	ACTR; AIB 1; AIB-1; AIB1; Amplified in breast cancer 1; Amplified in breast cancer 1 protein; bHLHe42; CAGH 16; CAGH16; CBP interacting protein; CBP-interacting protein; Class E basic helix-loop-helix protein 42; CTG 26; CTG26; KAT13B / AIB1; MGC141848; NCOA 3; NCOA 3; NCoA-3; Ncoa3; NCOA3_HUMAN; Nuclear receptor coactivator 3; pCIP; RAC 3; RAC-3; RAC3; Receptor associated coactivator 3; Receptor-associated coactivator 3; SRC 3; SRC-3; SRC3; Steroid receptor coactivator protein 3; Thyroid hormone receptor activator molecule 1; TNRC14; TNRC16; TRAM 1; TRAM-1; TRAM1.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Pig,Horse,Rabbit,Guinea Pig,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	155kDa
Cellular localization:	cytoplasmicThe cell membrane
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated Synthesised phosphopeptide derived from human SRC-3 around the phosphorylation site of Thr24:D(p-T)PG
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of

	antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	<p>KAT13B/AIB1 interacts in a ligand-dependent fashion with the estrogen receptor to enhance estrogen-dependent transcriptional gene activation. The protein encoded by this gene is a nuclear receptor coactivator that interacts with nuclear hormone receptors to enhance their transcriptional activator functions. The encoded protein has histone acetyltransferase activity and recruits KAT3A / KAT3B as part of a multisubunit coactivation complex. This protein is initially found in the cytoplasm but is translocated into the nucleus upon phosphorylation. Two transcript variants encoding different isoforms have been found for this gene. In addition, a polymorphic repeat region is found in the C-terminus of the encoded protein.</p> <p>Function: Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone-dependent fashion. Plays a central role in creating a multisubunit coactivator complex, which probably acts via remodeling of chromatin. Involved in the coactivation of different nuclear receptors, such as for steroids (GR and ER), retinoids (RARs and RXRs), thyroid hormone (TRs), vitamin D3 (VDR) and prostanoids (PPARs). Displays histone acetyltransferase activity. Also involved in the coactivation of the NF-kappa-B pathway via its interaction with the NFKB1 subunit. Interacts with PSMB9.</p> <p>Subunit: Interacts with CARM1. Present in a complex containing NCOA2, IKKA, IKKB, IKBKG and the histone acetyltransferase protein CREBBP. Interacts with CASP8AP2, NR3C1 and PCAF. Interacts with ATAD2 and this interaction is enhanced by estradiol.</p> <p>Subcellular Location: Cytoplasm. Nucleus. Note=Mainly cytoplasmic and weakly nuclear. Upon TNF activation and subsequent phosphorylation, it translocates from the cytoplasm to the nucleus.</p> <p>Tissue Specificity: Widely expressed. High expression in heart, skeletal muscle, pancreas and placenta. Low expression in brain, and very low in lung, liver and kidney.</p> <p>Post-translational modifications: Acetylated by CREBBP. Acetylation occurs in the RID domain, and disrupts the interaction with nuclear receptors and regulates its function. Methylated by CARM1 (By similarity). Phosphorylated by IKK complex. Regulated its function. Phosphorylation at Ser-601 by CK1 promotes coactivator function.</p> <p>Similarity: Belongs to the SRC/p160 nuclear receptor coactivator family. Contains 1 basic helix-loop-helix (bHLH) domain.</p>

Contains 1 PAS (PER-ARNT-SIM) domain.

SWISS:
Q9Y6Q9

Gene ID:
8202

Database links:

[Entrez Gene: 8202](#)Human

[Entrez Gene: 17979](#)Mouse

[Entrez Gene: 100135659](#)Pig

[Omin: 601937](#)Human

[SwissProt: Q9Y6Q9](#)Human

[SwissProt: O09000](#)Mouse


[Unigene: 592142](#)Human

[Unigene: 476883](#)Mouse

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Picture:



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-SRC3 (Thr24)) Polyclonal Antibody, Unconjugated (SL3408R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

www.sunlongbiotech.com