



Rabbit Anti-Phospho-MAP3K8 (Thr290) antibody

SL3455R

Product Name:	Phospho-MAP3K8 (Thr290)
Chinese Name:	磷酸化丝裂原活化蛋白激酶激酶8抗体
Alias:	MAP3K8 (Phospho T290); MAP3K8 (Phospho Thr290); P-MAP3K8/Tpl2 (Thr290); Phospho-Tpl2 (Thr290); P-Tpl2 (Thr290);c COT; Cancer Osaka thyroid oncogene; Cancer Osaka thyroid oncogene; CCOT; COT; COT proto oncogene serine/threonine protein kinase; EST; ESTF; Ewing sarcoma transformant; FLJ10486; M3K8_HUMAN; MAP3K 8; MAP3K8; Mitogen activated protein kinase kinase kinase 8; Mitogen-activated protein kinase kinase kinase 8; Proto oncogene cCot; Proto-oncogene c-Cot; Serine/threonine protein kinase cot; Serine/threonine-protein kinase cot; TPL 2; TPL-2; TPL2; Tumor progression locus 2.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Chicken,Dog,Pig,Cow,Horse,Rabbit,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	53kDa
Cellular localization:	cytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated Synthesised phosphopeptide derived from human MAP3K8/Tpl2 around the phosphorylation site of Thr290:RG(p-T)EI
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of


	antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	<p>This gene is an oncogene that encodes a member of the serine/threonine protein kinase family. The encoded protein localizes to the cytoplasm and can activate both the MAP kinase and JNK kinase pathways. This protein was shown to activate IkappaB kinases, and thus induce the nuclear production of NF-kappaB. This protein was also found to promote the production of TNF-alpha and IL-2 during T lymphocyte activation. This gene may also utilize a downstream in-frame translation start codon, and thus produce an isoform containing a shorter N-terminus. The shorter isoform has been shown to display weaker transforming activity. Alternate splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq, Sep 2011].</p> <p>Function: Required for TLR4 activation of the MEK/ERK pathway. Able to activate NF-kappa-B 1 by stimulating proteasome-mediated proteolysis of NF-kappa-B 1/p105. Plays a role in the cell cycle. The longer form has some transforming activity, although it is much weaker than the activated cot oncoprotein.</p> <p>Subunit: Forms a ternary complex with NFKB1 and TNIP2.</p> <p>Subcellular Location: Cytoplasm.</p> <p>Tissue Specificity: Expressed in several normal tissues and human tumor-derived cell lines.</p> <p>Post-translational modifications: Autophosphorylated. Isoform 1 undergoes phosphorylation mainly on Ser residues, and isoform 2 on both Ser and Thr residues.</p> <p>Similarity: Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily. Contains 1 protein kinase domain.</p> <p>SWISS: P41279</p> <p>Gene ID: 1326</p> <p>Database links: Entrez Gene: 1326Human Entrez Gene: 26410Mouse</p>

[Entrez Gene: 116596](#)Rat
[Oimim: 191195](#)Human
[SwissProt: P41279](#)Human
[SwissProt: Q07174](#)Mouse
[SwissProt: Q63562](#)Rat
[Unigene: 432453](#)Human
[Unigene: 3275](#)Mouse
[Unigene: 9939](#)Rat

Important Note:


This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Picture:



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-MAP3K8 (Thr290)) Polyclonal Antibody, Unconjugated (SL3455R) at 1:200 overnight at 4°C, followed by

operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: human colon carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-Phospho-MAP3K8/Tpl2(Thr290) Polyclonal Antibody,

Unconjugated(SL3455R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining