



## Rabbit Anti-phospho-CBL2 (Tyr674) antibody

SL4071R

<b>Product Name:</b>	phospho-CBL2 (Tyr674)
<b>Chinese Name:</b>	磷酸化原癌基因CBL2抗体
<b>Alias:</b>	phospho-C CBL (Tyr674); CBL2 (phospho-Tyr674); CBL2 (phospho-Y674); p-CBL2 (Tyr674); p-CBL2 (Y674); C CBL; Cas Br M (murine) ecotropic retroviral transforming sequence; Cas-Br-M (murine) ecotropic retroviral transforming sequence; Casitas B lineage lymphoma proto oncogene; Casitas B-lineage lymphoma proto-oncogene; CBL 2; cbl; CBL_HUMAN; CBL2; E3 ubiquitin protein ligase CBL; E3 ubiquitin-protein ligase CBL; Oncogene CBL2; Proto oncogene c CBL; Proto-oncogene c-CBL; RGD1561386; RING finger protein 55; RNF55; Signal transduction protein CBL.
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human,Mouse,Rat,Chicken,Dog,Pig,Cow,Horse,Rabbit,
<b>Applications:</b>	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	100kDa
<b>Cellular localization:</b>	cytoplasmicThe cell membrane
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated Synthesised phosphopeptide derived from human C CBL around the phosphorylation site of Tyr674:AI(p-Y)SL
<b>Isotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	<p>Participates in signal transduction in hematopoietic cells. Adapter protein that functions as a negative regulator of many signaling pathways that start from receptors at the cell surface. Acts as an E3 ubiquitin-protein ligase, which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome. Recognizes activated receptor tyrosine kinases, including PDGFA, EGF and CSF1, and terminates signaling.</p> <p><b>Function:</b>  Adapter protein that functions as a negative regulator of many signaling pathways that are triggered by activation of cell surface receptors. Acts as an E3 ubiquitin-protein ligase, which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome. Recognizes activated receptor tyrosine kinases, including KIT, FLT1, FGFR1, FGFR2, PDGFRA, PDGFRB, EGFR, CSF1R, EPHA8 and KDR and terminates signaling. Recognizes membrane-bound HCK and other kinases of the SRC family and mediates their ubiquitination and degradation. Participates in signal transduction in hematopoietic cells. Plays an important role in the regulation of osteoblast differentiation and apoptosis. Essential for osteoclastic bone resorption. The Tyr-731 phosphorylated form induces the activation and recruitment of phosphatidylinositol 3-kinase to the cell membrane in a signaling pathway that is critical for osteoclast function.</p> <p><b>Subunit:</b>  Interacts (phosphorylated at Tyr-731) with PIK3R1. Associates with NCK via its SH3 domain. The phosphorylated C-terminus interacts with CD2AP via its second SH3 domain. Binds to UBE2L3. Interacts with adapters SLA, SLA2 and with the phosphorylated C-terminus of SH2B2. Interacts with EGFR, SYK and ZAP70 via the highly conserved Cbl-N region. Also interacts with SORBS1 and INPPL1/SHIP2. Interacts with phosphorylated LAT2. May interact with CBLB (By similarity). Interacts with ALK, AXL, BLK, FGR and FGFR2. Interacts with CSF1R, EPHB1, FLT1, KDR, PDGFRA and PDGFRB; regulates receptor degradation through ubiquitination. Interacts with HCK and LYN. Interacts with TEK/TIE2 (tyrosine phosphorylated).</p> <p><b>Subcellular Location:</b>  Cytoplasm. Cell membrane. Note=Colocalizes with FGFR2 in lipid rafts at the cell membrane.</p> <p><b>Post-translational modifications:</b>  Phosphorylated on tyrosine residues by ALK, EGFR, SYK, FYN and ZAP70 (By similarity). Phosphorylated on tyrosine residues in response to FLT1 and KIT signaling. Phosphorylated on tyrosine residues by INSR and FGR. Phosphorylated on several tyrosine residues by constitutively activated FGFR3. Not phosphorylated at Tyr-731 by FGFR3. Phosphorylated on tyrosine residues by activated CSF1R, PDGFRA and PDGFRB. Phosphorylated on tyrosine residues by HCK. Ubiquitinated, leading to its degradation via the proteasome.</p>

**DISEASE:**

Defects in CBL are the cause of Noonan syndrome-like disorder with or without juvenile myelomonocytic leukemia (NSLL) [MIM:613563]. A syndrome characterized by a phenotype reminiscent of Noonan syndrome. Clinical features are highly variable, including facial dysmorphism, short neck, developmental delay, hyperextensible joints and thorax abnormalities with widely spaced nipples. The facial features consist of triangular face with hypertelorism, large low-set ears, ptosis, and flat nasal bridge. Some patients manifest cardiac defects.

**Similarity:**

Contains 1 Cbl-PTB (Cbl-type phosphotyrosine-binding) domain.

Contains 1 RING-type zinc finger.

Contains 1 UBA domain.

**SWISS:**

P22681

**Gene ID:**

867

**Database links:**

[Entrez Gene: Human](#)

[Entrez Gene: 12402](#) Mouse

[Entrez Gene: 500985](#) Rat

[Omim: 165360](#) Human

[SwissProt: P22681](#) Human

[SwissProt: P22682](#) Mouse

[Unigene: 504096](#) Human

[Unigene: 266871](#) Mouse

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

C CBL是Ubiquitin-蛋白酶体(ubiquitin proteasome)通路中的一个新的RING Finger型Ubiquitin连接酶(ubiquitinligase ,E3).C CBL可以介导受体酪氨酸激酶和非受体酪氨酸受体激酶的降解.

