



Rabbit Anti-ASGPR1/FITC Conjugated antibody

SL4119R-FITC

Product Name:	Anti-ASGPR1/FITC
Chinese Name:	FITC标记的唾液酸glycoprotein受体1抗体
Alias:	ASGR; Asgr1; ASGPR1; Asgr; Asgr-1; MGC108731; RATRHL1; RHL1; ASGR1_HUMAN; Asialoglycoprotein receptor 1; ASGP-R 1; ASGPR 1; C-type lectin domain family 4 member H1; Hepatic lectin H1; HL-1.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Pig,Cow,Horse,Rabbit,
Applications:	not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	32kDa
Cellular localization:	The cell membrane
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human ASGPR1
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Product Detail:	background: ASGR is a heterooligomeric receptor that is abundantly expressed on the sinusoidal surface of the hepatic plasma membrane. It is an endocytic receptor that rapidly binds and internalizes galactose-terminated glycoproteins (asialoglycoproteins or ASGP) from the circulation. The mouse ASGPR belongs to the long-form subfamily of the C-type/Ca ²⁺ dependent lectin family. It is a complex of two noncovalently-linked and highly homologous subunits, a major 42 kDa glycoprotein ASGPR1(MHL-1) and a

minor 51 kDa glycoprotein ASGR2 (MHL-2). ASGPR1 is synthesized as a type II transmembrane protein that contains a cytosolic N-terminal domain, a single transmembrane segment, and an extracellular domain which contains two important structural regions. The first is a stalk domain that contributes to noncovalent oligomerization, and the second is a Ca²⁺-dependent carbohydrate binding domain at the very C-terminus that is unusually stabilized by three ions. The aa sequence of mouse ASGPR1 ECD is 89% and 79% identical to the ASGPR1 ECD of rat and human, respectively.

Function:

Mediates the endocytosis of plasma glycoproteins to which the terminal sialic acid residue on their complex carbohydrate moieties has been removed. The receptor recognizes terminal galactose and N-acetylgalactosamine units. After ligand binding to the receptor, the resulting complex is internalized and transported to a sorting organelle, where receptor and ligand are disassociated. The receptor then returns to the cell membrane surface.

Subunit:

Interacts with LASS2.

Subcellular Location:

Membrane; Single-pass type II membrane protein.

Tissue Specificity:

Expressed exclusively in hepatic parenchymal cells.

Post-translational modifications:

Phosphorylated on a cytoplasmic Ser residue.

Similarity:

Contains 1 C-type lectin domain.

Database links:

[Entrez Gene: 509121](#)Cow

[Entrez Gene: 432](#)Human

[Entrez Gene: 11889](#)Mouse

[Entrez Gene: 24210](#)Rat

[Omim: 108360](#)Human

[SwissProt: P07306](#)Human

[SwissProt: P34927](#)Mouse

[SwissProt: P02706](#)Rat

[Unigene: 12056](#)Human

[Unigene: 6559](#)Mouse

[Unigene: 44300](#)Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

www.sunlongbiotech.com