



## Rabbit Anti-ZMYND17 antibody

SL4395R

<b>Product Name:</b>	ZMYND17
<b>Chinese Name:</b>	ZMYND17蛋白抗体
<b>Alias:</b>	FLJ39565; Putative protein MSS51 homolog, mitochondrial; Zinc finger MYND domain containing 17; Zinc finger MYND domain containing protein 17; Zinc finger MYND type containing 17; ZMYND 17.
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human,
<b>Applications:</b>	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	51kDa
<b>Cellular localization:</b>	The nucleus
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated synthetic peptide derived from human ZMYND17:361-460/460
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	Preservative: 15mM Sodium Azide, Constituents: 1% BSA, 0.01M PBS, pH 7.4
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	ZMYND17 contains 1 MYND-type zinc finger. The exact function of ZMYND17 remains unknown.  SWISS: Q4VC12

**Gene ID:**  
118490

**Database links:**

[Entrez Gene: 118490](#) Human

[Omim: 614773](#) Human


[SwissProt: Q4VC12](#) Human

[Unigene: 631823](#) Human

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**Picture:**



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZMYND17) Polyclonal

	Antibody, Unconjugated (SL4395R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.
--	--

[www.sunlongbiotech.com](http://www.sunlongbiotech.com)