

Rabbit Anti-PTPNS1L2 antibody

SL4432R

PTPNS1L2
信号调节蛋白SIRPD抗体
Protein tyrosine phosphatase non receptor type substrate 1 like 2; PTPNS1L2; Signal regulatory protein delta; SIRPD.
Rabbit
Polyclonal
Human,
WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100- 500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
22kDa
Secretory protein
Lyophilized or Liquid
1mg/ml
KLH conjugated synthetic peptide derived from human PTPNS1L2:81-180/197
IgG
affinity purified by Protein A
Preservative: 15mM Sodium Azide, Constituents: 1% BSA, 0.01M PBS, pH 7.4
Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed
 SIRPD (Signal Regulatory Protein Delta) is a Protein Coding gene. An important paralog of this gene is SIRPG. Similarity: Contains 1 Ig-like V-type (immunoglobulin-like) domain.

	SWISS:
	Q9H106
	C ID
	Gene ID:
	128646
	Database links:
	Entrez Gene: 128646 Human
	SwissProt: Q9H106 Human
	Unigene: 729560 Human
	Important Note:
	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Picture:	
	Paraformaldehyde-fixed, paraffin embedded (Human kidney tissue); Antigen
	retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous
	peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat
	serum) at 37°C for 30min; Antibody incubation with (PTPNS1L2) Polyclonal

Antibody, Unconjugated (SL4432R) at 1:400 overnight at 4°C, followed by
operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

www.sunonobiotech.com