



## Rabbit Anti-phospho-MAPKAPK5 (Thr182) antibody

SL5504R

<b>Product Name:</b>	phospho-MAPKAPK5 (Thr182)
<b>Chinese Name:</b>	磷酸化p38调节/激活蛋白激酶抗体
<b>Alias:</b>	p-PRAK(phospho Thr182); MAPKAPK5(phospho T182); MAP kinase-activated protein kinase 5; MAPKAP kinase 5; MAPKAPK5; mitogen-activated protein kinase-activated protein kinase 5; p38-regulated/activated protein kinase.
<b>Organism Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>React Species:</b>	Human,Mouse,Rat,Chicken,Dog,Pig,Cow,Horse,Rabbit,
<b>Applications:</b>	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight:</b>	52kDa
<b>Cellular localization:</b>	The nucleuscytoplasmic
<b>Form:</b>	Lyophilized or Liquid
<b>Concentration:</b>	1mg/ml
<b>immunogen:</b>	KLH conjugated Synthesised phosphopeptide derived from human MAPKAPK5 around the phosphorylation site of Thr182:LM(p-T)PQ
<b>Lsotype:</b>	IgG
<b>Purification:</b>	affinity purified by Protein A
<b>Storage Buffer:</b>	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
<b>Storage:</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>PubMed:</b>	<a href="#">PubMed</a>
<b>Product Detail:</b>	The serine/threonine kinase PRAK is activated in response to cellular stress and

proinflammatory cytokines, through its phosphorylation by MAP kinases including MAPK1/ERK, MAPK14/p38 alpha, and MAPK11/p38 beta. PRAK has been reported to have a putative tumor suppressor function by mediating senescence upon activation by p38 in response to oncogenic ras. It is thought that phosphorylation of p53 by PRAK following activation of p38 MAPK by ras plays an important role in ras induced senescence and tumor suppression.

**Function:**

Tumor suppressor serine/threonine-protein kinase involved in mTORC1 signaling and post-transcriptional regulation. Phosphorylates FOXO3, ERK3/MAPK6, ERK4/MAPK4, HSP27/HSPB1, p53/TP53 and RHEB. Acts as a tumor suppressor by mediating Ras-induced senescence and phosphorylating p53/TP53. Involved in post-transcriptional regulation of MYC by mediating phosphorylation of FOXO3: phosphorylation of FOXO3 leads to promote nuclear localization of FOXO3, enabling expression of miR-34b and miR-34c, 2 post-transcriptional regulators of MYC that bind to the 3'UTR of MYC transcript and prevent MYC translation. Acts as a negative regulator of mTORC1 signaling by mediating phosphorylation and inhibition of RHEB. Part of the atypical MAPK signaling via its interaction with ERK3/MAPK6 or ERK4/MAPK4: the precise role of the complex formed with ERK3/MAPK6 or ERK4/MAPK4 is still unclear, but the complex follows a complex set of phosphorylation events: upon interaction with atypical MAPK (ERK3/MAPK6 or ERK4/MAPK4), ERK3/MAPK6 (or ERK4/MAPK4) is phosphorylated and then mediates phosphorylation and activation of MAPKAPK5, which in turn phosphorylates ERK3/MAPK6 (or ERK4/MAPK4). Mediates phosphorylation of HSP27/HSPB1 in response to PKA/PRKACA stimulation, inducing F-actin rearrangement.

**Subunit:**

Interacts with ERK3/MAPK6 and ERK4/MAPK4 (via FRIEDE motif); the interaction is direct. Interacts with YWHAE; the interaction prevents phosphorylation of HSP27/HSPB1 leading to disrupt F-actin polymerization. Interacts with SQSTM1.

**Subcellular Location:**

Cytoplasm. Nucleus.

**Tissue Specificity:**

Expressed ubiquitously.

**Post-translational modifications:**

Phosphorylated on Thr-182 ERK3/MAPK6 or ERK4/MAPK4; which is the regulatory phosphorylation site and is located on the T-loop/loop 12, leading to activation. Phosphorylation at Thr-182 by p38-alpha/MAPK14, p38-beta/MAPK11 is subject to debate. Phosphorylated at Ser-115 by PKA/PRKACA, leading to localization to the cytoplasm. Autophosphorylated (By similarity).

**Similarity:**

Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family.

Contains 1 protein kinase domain.

**SWISS:**  
Q8IW41

**Gene ID:**  
8550

**Database links:**

[Entrez Gene: 8550](#)Human

[Entrez Gene: 17165](#)Mouse

[Entrez Gene: 100153404](#)Pig

[Entrez Gene: 498183](#)Rat

[Olim: 606723](#)Human

[SwissProt: Q8IW41](#)Human

[SwissProt: O54992](#)Mouse


[Unigene: 413901](#)Human

[Unigene: 272206](#)Mouse

[Unigene: 17463](#)Rat


**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



**Picture:**

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-MAPKAPK5 (Thr182)) Polyclonal Antibody, Unconjugated (SL5504R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-MAPKAPK5(Thr182)) Polyclonal Antibody, Unconjugated (SL5504R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.