



Rabbit Anti-MOCOS antibody

SL5936R

Product Name:	MOCOS
Chinese Name:	原癌基因丝氨酸/苏氨酸蛋白激酶MOS抗体
Alias:	c mos; c-mos; MSV; y Oocyte maturation factor mos; Proto oncogene serine / threonine protein kinase mos; FLJ20733; HMCS; MCS; MoCo sulfurase; Molybdenum cofactor sulfurase; MOS; v mos Moloney murine sarcoma viral oncogene homolog; MOCOS_HUMAN.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Dog,Pig,Horse,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	98kDa
Cellular localization:	cytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human MOCOS:551-650/888
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	MOCOS sulfurates the molybdenum cofactor of xanthine dehydrogenase (XDH; MIM 607633) and aldehyde oxidase (AOX1; MIM 602841), which is required for their enzymatic activities (Ichida et al., 2001 [PubMed 11302742]).[supplied by OMIM, Feb 2010]

Function:

Sulfurates the molybdenum cofactor. Sulfation of molybdenum is essential for xanthine dehydrogenase (XDH) and aldehyde oxidase (ADO) enzymes in which molybdenum cofactor is liganded by 1 oxygen and 1 sulfur atom in active form.

DISEASE:

Defects in MOCOS are the cause of xanthinuria type 2 (XU2) [MIM:603592]. Xanthinuria is characterized by excretion of very large amounts of xanthine in the urine and a tendency to form xanthine stones. Uric acid is strikingly diminished in serum and urine. In addition, patient suffering of XU2 cannot metabolize allopurinol into oxypurinol due to dual deficiency of xanthine dehydrogenase and aldehyde oxidase.

Similarity:

Belongs to the class-V pyridoxal-phosphate-dependent aminotransferase family. MOCOS subfamily. Contains 1 MOSC domain.

SWISS:

Q96EN8

Gene ID:

55034

Database links:

[Entrez Gene: 55034](#) Human

[Omim: 613274](#) Human

[SwissProt: Q96EN8](#) Human

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.