



Rabbit Anti-G21 protein/NPR2L antibody

SL6079R

Product Name:	G21 protein/NPR2L
Chinese Name:	G21蛋白质抗体
Alias:	G21 protein; Gene 21 protein; Homologous to yeast nitrogen permease (candidate tumor suppressor); Homologous to yeast nitrogen permease; NPR 2L; NPR L2; NPR like 2; NPR2 like; NPRL 2; NPRL2; Tumor suppressor candidate 4; TUSC 4; TUSC 4 protein; TUSC4; TUSC4 protein; NPRL2_HUMAN.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Pig,Cow,Horse,Rabbit,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	44kDa
Cellular localization:	cytoplasmicThe cell membrane
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human NPR2L:221-320/380
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	NPR2L is homologous to yeast nitrogen permease and is a candidate tumor suppressor, being a negative regulator of cell cycle. Most abundant in skeletal muscle, followed by brain, liver, and pancreas, with lower amounts in lung, kidney, placenta, and heart. Expressed in most lung cancer cell lines tested. There are two isoforms, produced by

alternative splicing.

Function:

Suppresses Src-dependent tyrosine phosphorylation and activation of PDPK1 and its downstream signaling. Down-regulates PDPK1 kinase activity by interfering with tyrosine phosphorylation at the Tyr-9 Tyr-373 and Tyr-376 residues. May act as a tumor suppressor. Suppresses cell growth and enhanced sensitivity to various anticancer drugs.

Subunit:

Forms a heterodimer with NPRL3. Interacts with PDPK1.

Tissue Specificity:

Most abundant in skeletal muscle, followed by brain, liver and pancreas, with lower amounts in lung, kidney, placenta and heart. Expressed in most lung cancer cell lines tested.

Similarity:

Belongs to the NPR2 family.

SWISS:

Q8WTW4

Gene ID:

10641

Database links:

[Entrez Gene: 508487](#) Cow

[Entrez Gene: 10641](#) Human

[Entrez Gene: 56032](#) Mouse

[Omim: 607072](#) Human

[SwissProt: Q5E9U9](#) Cow

[SwissProt: Q8WTW4](#) Human


[SwissProt: Q9WUE4](#) Mouse

[Unigene: 437083](#) Human

[Unigene: 11337](#) Mouse

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



Picture:

Sample:


HepG2(Human) Cell Lysate at 30 ug

Primary: Anti-NPR2L (SL6079R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 44 kD

Observed band size: 44 kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (G21 protein/NPR2L) Polyclonal Antibody, Unconjugated (SL6079R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.