

# Rabbit Anti-GPR48 antibody

# SL6676R

<b>Product Name:</b>	GPR48			
Chinese Name:	G protein-coupled receptor48抗体			
Alias:	G protein coupled receptor 48; G-protein coupled receptor 48; GPR48; GPCR GPR48; Leucine-rich repeat-containing G protein-coupled receptor 4; Leucine-rich repeat-containing G-protein coupled receptor 4; LGR4; LGR4_HUMAN.			
Organism Species:	Rabbit			
Clonality:	Polyclonal			
React Species:	Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Rabbit,			
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.			
Molecular weight:	104kDa			
Cellular localization:	The cell membrane			
Form:	Lyophilized or Liquid			
Concentration:	lmg/ml			
immunogen:	KLH conjugated synthetic peptide derived from human LGR4:65-170/951 <extracellular></extracellular>			
Lsotype:	IgG			
Purification:	affinity purified by Protein A			
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.			
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.			
PubMed:	<u>PubMed</u>			
Product Detail:	G protein-coupled receptors (GPCRs), also designated seven transmembrane (7TM) receptors or heptahelical receptors, interact with G proteins (heterotrimeric GTPases) to synthesize intracellular second messengers, such as diacylglycerol, cyclic AMP, inositol phosphates and calcium ions. Their diverse biological functions range from vision and			

olfaction to neuronal and endocrine signaling and are involved in many pathological conditions. LGR4 (leucine-rich repeat-containing G protein-coupled receptor 4), also known as GPR48, is a 951 amino acid multi-pass membrane protein that contains 15 LRR (leucine-rich repeats) and belongs to the GPCR family. Expressed in multiple tissues, including testis, ovary, placenta, stomach, heart, kidney, pancreas and spleen, LGR4 functions as an orphan receptor that may be involved in physiologic activities throughout the cell. LGR4 is overexpressed in various cancer types and is thought to enhance carcinoma invasiveness and metastasis, suggesting an important role in tumor progression.

#### **Function:**

Orphan receptor.

#### **Subcellular Location:**

Cell membrane; Multi-pass membrane protein.

### Tissue Specificity:

Expressed in multiple steroidogenic tissues: placenta, ovary, testis and adrenal. Expressed also in spinal cord, thyroid, stomach, trachea, heart, pancreas, kidney, prostate and spleen.

### Similarity:

Belongs to the G-protein coupled receptor 1 family.

Contains 15 LRR (leucine-rich) repeats.

Contains 1 LRRNT domain.

## **SWISS:**

O9BXB1

#### Gene ID:

55366

#### Database links:

Entrez Gene: 55366Human

Entrez Gene: 107515 Mouse

Entrez Gene: 286994Rat

Omim: 606666Human

SwissProt: Q9BXB1Human

SwissProt: A2ARI4Mouse

SwissProt: Q9Z2H4Rat

Unigene: 502176Human


Uni	gene:	27735	1 Mouse
	_		

Unigene: 480424 Mouse

Unigene: 162671Rat

## Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



## Picture:

Sample: Hela Cell (Human) Lysate at 40 ug

Primary: Anti-GPR48 (SL6676R) at 1/300 dilution

Secondary: HRP conjugated Goat-Anti-rabbit IgG (SL6676R) at 1/5000 dilution

Predicted band size: 104 kD

Observed band size: 104 kD