



Rabbit Anti-ZNF137 antibody

SL7153R

| | |
|-------------------------------|---|
| Product Name: | ZNF137 |
| Chinese Name: | Zinc finger protein137抗体 |
| Alias: | pHZ 30; ZN137_HUMAN; Putative zinc finger protein 137; zinc finger protein 137, pseudogene; ZNF137P. |
| Organism Species: | Rabbit |
| Clonality: | Polyclonal |
| React Species: | Human, |
| Applications: | IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user. |
| Molecular weight: | 24kDa |
| Cellular localization: | The nucleus |
| Form: | Lyophilized or Liquid |
| Concentration: | 1mg/ml |
| immunogen: | KLH conjugated synthetic peptide derived from human ZNF137:1-100/207 |
| Lsotype: | IgG |
| Purification: | affinity purified by Protein A |
| Storage Buffer: | 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol. |
| Storage: | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |
| PubMed: | PubMed |
| Product Detail: | May be involved in transcriptional regulation. Function: May be involved in transcriptional regulation. Subcellular Location: |

Nucleus (Potential).

Similarity:

Belongs to the krueppel C2H2-type zinc-finger protein family.
Contains 5 C2H2-type zinc fingers.

SWISS:

P52743

Gene ID:

7696

Database links:

[Entrez Gene: 7696](#) Human


[SwissProt: P52743](#) Human

[Unigene: 373648](#) Human

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Picture:



Paraformaldehyde-fixed, paraffin embedded (human gastric); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by

3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF137) Polyclonal Antibody, Unconjugated (SL7153R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

www.sunlongbiotech.com