



Rabbit Anti-SP3 antibody

SL7342R

Product Name:	SP3
Chinese Name:	转录因子SP3抗体
Alias:	D130027J01Rik; DKFZp686O1631; GC binding transcription factor Sp 3; GC binding transcription factor Sp3; MGC105187; OTTMUSP00000014207; SP 3; Sp 3 transcription factor; SP3; Sp3 transcription factor; SP3_HUMAN; Specificity protein 3; SPR 2; SPR-2; SPR2; Transcription factor SP 3; Transcription factor SP3; Transcription factor Sp3.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Chicken,Cow,Rabbit,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	82kDa
Cellular localization:	The nucleus
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human SP3:651-750/781
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	The Sp transcription factor family includes Sp1, Sp2, Sp3 (SPR-2) and Sp4 (SPR-1). Sp transcription factors share similar structures but do not share simi-lar functions. All four proteins contain a highly conserved DNA-binding domain composed of three zinc

fingers at the C-terminus. Sp family members bind the consensus sequence GGGGCGGGGC and other closely related sequences which are known as GC boxes. Sp1, Sp3 and Sp4 share a high affinity for GC boxes while Sp2 does not. Sp2 only weakly binds to GT boxes. Sp1, Sp2 and Sp3 are ubiquitously expressed, while Sp4 is abundantly expressed in brain with limited expression in other tissues. Sp1 and Sp3, but not Sp2 or Sp4, interact with E2, a regulatory element for the β 4 subunit of neuronal nicotinic acetylcholine receptors. Sp3 is the only Sp member to inhibit Sp1 and Sp4

Function:

Transcriptional factor that can act as an activator or repressor depending on isoform and/or post-translational modifications. Binds to GT and GC boxes promoter elements. Competes with SP1 for the GC-box promoters. Weak activator of transcription but can activate a number of genes involved in different processes such as cell-cycle regulation, hormone-induction and house-keeping.

Subunit:

Interacts with HLTF; the interaction may be required for basal transcriptional activity of HLTF. Interacts with HDAC1; the interaction deacetylates SP3 and regulates its transcriptional activity. Interacts with HDAC2 (preferably the CK2-phosphorylated form); the interaction deacetylates SP3 and regulates its transcriptional activity. Interacts with MEIS2 isoform 4 and PBX1 isoform PBX1a.

Subcellular Location:

Nucleus. Nucleus > PML body. Localizes to the nuclear periphery and in nuclear dots when sumoylated. Some localization in PML nuclear bodies.

Tissue Specificity:

Ubiquitously expressed.

Post-translational modifications:

Not glycosylated.

Acetylated by histone acetyltransferase p300, deacetylated by HDACs.

Acetylation/deacetylation states regulate transcriptional activity. Acetylation appears to activate transcription. Alternate sumoylation and acetylation at Lys-551 also control transcriptional activity. Ceramides can also regulate acetylation/deacetylation events through altering the interaction of HDAC with SP3. In vitro, C(18)-ceramides, but not C(16)-ceramides, increase the interaction of HDAC1 with SP3 and enhance the deacetylation of SP3 and the subsequent repression of the TERT promoter.

Sumoylated on all isoforms. Sumoylated on 2 sites in longer isoforms with Lys-551 being the major site. Sumoylation at this site promotes nuclear localization to the nuclear periphery, nuclear dots and PML nuclear bodies. Sumoylation on Lys-551 represses the transactivation activity, except for the largest isoform, L-Sp3, which has little effect on transactivation. Alternate sumoylation and acetylation at Lys-551 also control transcriptional activity.

Similarity:

Belongs to the Sp1 C2H2-type zinc-finger protein family.
Contains 3 C2H2-type zinc fingers.

SWISS:

Q02447

Gene ID:

6670

Database links:

[Entrez Gene: 6670](#) Human

[Entrez Gene: 20687](#) Mouse

[Entrez Gene: 367846](#) Rat

[Omim: 601804](#) Human

[SwissProt: Q02447](#) Human

[SwissProt: O70494](#) Mouse

[Unigene: 531587](#) Human

[Unigene: 124328](#) Mouse

[Unigene: 446209](#) Mouse

[Unigene: 465887](#) Mouse

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.