


## Rabbit Anti-Tankyrase 2 antibody

## SL7389R

Tankyrase 2
端粒调控因子2抗体
ADP-ribosyltransferase diphtheria toxin-like 6; ARTD6; PARP 5b; PARP 5c; PARP-5b; PARP-5c; PARP5B; PARP5C; pART6; Poly [ADP ribose] polymerase 5B; Poly [ADP-ribose] polymerase 5B; TANK2; Tankyrase 2; Tankyrase II; Tankyrase like protein; Tankyrase related protein; Tankyrase TRF1 interacting ankyrin related ADP ribose polymerase 2; Tankyrase-2; Tankyrase-like protein; Tankyrase-related protein; TNKL; TNKS 2; TNKS-2; TNKS2; TNKS2_HUMAN; TRF1 Interacting Ankyrin Related ADP ribose polymerase 2; TRF1-interacting ankyrin-related ADP-ribose polymerase 2.
Rabbit
Polyclonal
Human, Mouse, Rat, Dog, Pig, Cow, Horse,
ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
127kDa
cytoplasmic
Lyophilized or Liquid
lmg/ml
KLH conjugated synthetic peptide derived from human Tankyrase 2:1-100/1166
IgG
affinity purified by Protein A
0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed



## Picture:

Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Tankyrase 2) Polyclonal Antibody, Unconjugated (SL7389R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.