

Rabbit Anti-CCDC136 antibody

SL8090R

Product Name:	CCDC136
Chinese Name:	鼻咽癌相关蛋白6抗体
Alias:	CC136_HUMAN; Ccdc136; Coiled-coil domain-containing protein 136; DKFZP434G156; KIAA1793; NAG6; Nasopharyngeal carcinoma-associated gene 6 protein.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human, Mouse, Rat,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	134kDa
Cellular localization:	The cell membrane
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from Human CCDC136/NAG6:85-190/1154
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
	Expressed in gastric tissues. Down-regulated in gastric cancer.
Product Detail:	Subcellular Location: Membrane; Single-pass membrane protein (Potential).

Tissue Specificity:

Expressed in gastric tissues. Down-regulated in gastric cancer.

SWISS:

Q96JN2

Gene ID:


64753

Database links:

UniProtKB/Swiss-Prot: Q96JN2.3

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



Picture:

Tissue/cell: Rat stomach tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-CCDC136 Polyclonal Antibody, Unconjugated(SL8090R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and

DAB(C-0010) staining

www.surlondbiotech.com