


Rabbit Anti-CCDC160 antibody

SL8123R


CCDC160	
卷曲螺旋结构域蛋白160抗体	
CC160_HUMAN; CCDC160; Coiled-coil domain containing 160.	
Rabbit	
Polyclonal	
Human, Mouse, Rat, Cow, Sheep,	
ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:100-500 (Paraffin sections	
need antigen repair)	
not yet tested in other applications.	
optimal dilutions/concentrations should be determined by the end user.	
38kDa	
The nucleuscytoplasmic	
Lyophilized or Liquid	
lmg/ml	
KLH conjugated synthetic peptide derived from Human CCDC160:182-280/325	
IgG	
affinity purified by Protein A	
0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.	
Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of	
	antibody the antibody is stable for at least two weeks at 2-4 °C.
	<u>PubMed</u>



Picture:

Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-FAM113A Polyclonal Antibody, Unconjugated(SL8123R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (Mouse small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CCDC160) Polyclonal Antibody, Unconjugated (SL8123R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.