




Rabbit Anti-USP15 antibody


SL8752R

Product Name:	USP15
Chinese Name:	去Ubiquitin化酶15抗体
Alias:	Deubiquitinating enzyme 15; Deubiquitinating enzyme; EC 3 1 2 15; KIAA0529; MGC131982; MGC149838; MGC74854; Ubiquitin Carboxy terminal Hydrolase 15; Ubiquitin carboxyl-terminal hydrolase 15; Ubiquitin specific peptidase 15; Ubiquitin specific processing protease 15; Ubiquitin Specific Protease 15; Ubiquitin thioesterase 15; Ubiquitin thiolesterase 15; Ubiquitin-specific-processing protease 15; UBP 15; UBP15; UBP15 HUMAN; Unph 2; UNPH 4; Unph-2; Unph2; Unph4; USP 15.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Chicken,Dog,Pig,Cow,Horse,Rabbit,Sheep,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:100-500 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	112kDa
Cellular localization:	The nucleuscytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human USP15:601-700/981
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	



Picture:

Paraformaldehyde-fixed, paraffin embedded (Rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (USP15) Polyclonal Antibody, Unconjugated (SL8752R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (USP15) Polyclonal Antibody, Unconjugated (SL8752R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.