



Rabbit Anti-LRSAM1 antibody

SL9387R

Product Name:	LRSAM1
Chinese Name:	LRSAM1 蛋白抗体
Alias:	E3 ubiquitin protein ligase LRSAM1; hTAL; Leucine rich repeat and sterile alpha motif containing protein 1; RIFLE; TAL; Tsg101 associated ligase; LRSM1 HUMAN.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Cow,Horse,Sheep,
Applications:	WB=1:500-2000ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:50-200 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	84kDa
Cellular localization:	cytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated synthetic peptide derived from human LRSAM1:201-300/723
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
PubMed:	PubMed
Product Detail:	LRSAM1 is an E3 ubiquitin-protein ligase that mediates monoubiquitination of TSG101 at multiple sites, leading to inactivation of the ability of TSG101 to sort endocytic (EGF receptors) and exocytic (HIV-1 viral proteins) cargos. It selectively regulates cell adhesion molecules and plays a role in receptor endocytosis and viral budding. LRSAM1 contains a RING-type zinc finger, 5 leucine-rich repeats and 1 SAM (sterile alpha motif) domain. The coiled coil domains interact with the SB domain of TSG101.

The PTAP motifs mediate the binding to UEV domains. There are 3 isoforms produced by alternative splicing.

Function:

E3 ubiquitin-protein ligase that mediates monoubiquitination of TSG101 at multiple sites, leading to inactivate the ability of TSG101 to sort endocytic (EGF receptors) and exocytic (HIV-1 viral proteins) cargos.

Subunit:

Interacts with TSG101.

Subcellular Location:

Cytoplasm. Note=Displays a punctuate distribution and localizes to a submembranal ring.

Tissue Specificity:

Highly expressed in adult spinal cord motoneurons as well as in fetal spinal cord and muscle tissue.

DISEASE:

Defects in LRSAM1 are a cause of Charcot-Marie-Tooth disease type 2P (CMT2P) [MIM:614436]. CMT2P is an axonal form of Charcot-Marie-Tooth disease, a disorder of the peripheral nervous system, characterized by progressive weakness and atrophy, initially of the peroneal muscles and later of the distal muscles of the arms. Charcot-Marie-Tooth disease is classified in two main groups on the basis of electrophysiologic properties and histopathology: primary peripheral demyelinating neuropathies (designated CMT1 when they are dominantly inherited) and primary peripheral axonal neuropathies (CMT2). Neuropathies of the CMT2 group are characterized by signs of axonal degeneration in the absence of obvious myelin alterations, normal or slightly reduced nerve conduction velocities, and progressive distal muscle weakness and atrophy. Nerve conduction velocities are normal or slightly reduced.

Similarity:

Contains 6 LRR (leucine-rich) repeats.

Contains 1 RING-type zinc finger.

Contains 1 SAM (sterile alpha motif) domain.

SWISS:

Q6UWE0

Gene ID:

90678


Database links:

[Entrez Gene: 90678](#)Human

[Oimim: 610933](#)Human
[SwissProt: Q6UWE0](#)Human
[Unigene: 495188](#)Human

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



Picture:

Paraformaldehyde-fixed, paraffin embedded (mouse spinal cord); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LRSAM1) Polyclonal Antibody, Unconjugated (SL9387R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.