

# Rabbit Anti-PRR15 antibody

# SL9408R

Chinese Name:   富含脯氨酸蛋白15抗体	
OTTHUMP00000202286.  Organism Species: Rabbit Clonality: Polyclonal  React Species: Human,Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit,Sheep, ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:50-200 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight: 14kDa Cellular localization: The nucleuscytoplasmic Form: Lyophilized or Liquid Concentration: Img/ml	
Organism Species: Rabbit Clonality: Polyclonal React Species: Human,Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit,Sheep, ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:50-200 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight: Cellular localization: The nucleuscytoplasmic Form: Lyophilized or Liquid Concentration: Img/ml	
Clonality:  React Species:  Human,Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit,Sheep,  ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:50-200 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight: Cellular localization: The nucleuscytoplasmic Form: Lyophilized or Liquid Concentration: Img/ml	
React Species:  Human,Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit,Sheep,  ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:50- 200 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight: Cellular localization: The nucleuscytoplasmic Form: Lyophilized or Liquid Concentration: Img/ml	
Applications:  ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800ICC=1:100-500IF=1:50-200 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight: 14kDa  Cellular localization: The nucleuscytoplasmic  Form: Lyophilized or Liquid  Concentration: 1mg/ml	
Applications:  200 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight: 14kDa Cellular localization: The nucleuscytoplasmic  Form: Lyophilized or Liquid Concentration:  1mg/ml	
not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight: 14kDa Cellular localization: The nucleuscytoplasmic Form: Lyophilized or Liquid Concentration: 1mg/ml	
not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.  Molecular weight: 14kDa Cellular localization: The nucleuscytoplasmic Lyophilized or Liquid Concentration:  Img/ml	
Molecular weight:14kDaCellular localization:The nucleuscytoplasmicForm:Lyophilized or LiquidConcentration:1mg/ml	
Cellular localization:       The nucleuscytoplasmic         Form:       Lyophilized or Liquid         Concentration:       1mg/ml	
Form: Lyophilized or Liquid Concentration: lmg/ml	
Concentration: 1mg/ml	
3	
immunogon. KI H conjugated synthetic pentide derived from human DR P.15:41 120/120	
inimunogen. KEAT conjugated synthetic peptide derived from human I KK13.41-129/129	
Lsotype: IgG	
Purification: affinity purified by Protein A	
Storage Buffer: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.	
Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized	
Storage: antibody is stable at room temperature for at least one month and for greater than a	year
when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of	
antibody the antibody is stable for at least two weeks at 2-4 °C.	
PubMed: PubMed	
May have a role in proliferation and/or differentiation.	
Product Detail:	
May have a role in proliferation and/or differentiation (By similarity).	
Similarity:	

Belongs to the PRR15 family.

SWISS: Q8IV56

Gene ID: 222171

#### Database links:


Entrez Gene: 222171 Human

SwissProt: Q8IV56Human

Unigene: 728338Human

## **Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



### Picture:

Tissue/cell: A549 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PRR15) Polyclonal Antibody, Unconjugated (SL9408R) 1:100, 90

minutes at 37°C, followed by operating according to SP Kit(Rabbit) (sp-0023)
instructionsand DAB staining.

