

Basic xylanase (BAX) Activity Assay Kit

Note: Take two or three different samples for prediction before test.

Operation Equipment: Spectrophotometer

Catalog Number: AK0133

Size:50T/24S

Components:

Buffer: 50mL×1, storage at 4°C .

Reagent 1: 10mL×1, storage at 4°C .

Reagent 2: 15mL×1, storage at 4°C .

Standard: Powder×1, storage at 4°C . 10mg of xylose, add 0.667mL of distilled water to dissolve before use, prepare a 100μmol/ mL standard solution. Dilute 50 times to prepare 2μmol / mL xylose standard solution for use

Product Description:

Xylanase (EC 3.2. 1.8) is mainly produced by microorganisms and can catalyze the hydrolysis of xylan, also known as pentosanase or hemicellulase. It can decompose the cell wall of raw materials and β - glucan in brewing or feed industry. It is widely used in brewing and feed industry to reduce the viscosity of materials, promote the release of effective substances, reduce the non-starch polysaccharides in feeding, and promote the absorption and utilization of nutrients. In general, alkaline xylanase (Bax) is isolated from microorganisms with optimal growth pH of 9- 11.

BAX catalyzes the degradation of xylan into reducing oligosaccharides and monosaccharides in an alkaline environment, and further develops a color reaction with 3,5-dinitrosalicylic acid in a boiling water bath. The color of the reaction solution is proportional to the amount of reducing sugar produced by the enzymatic hydrolysis. The BAX activity can be calculated by measuring the increase rate of the absorbance of the reaction solution at 540nm.

Reagents and Equipment Required but Not Provided:

Spectrophotometer, adjustable transferpettor, balance, mortar/homogenizer, centrifuge, 1mL glass cuvette and distilled water.

Sample preparation:

1. Plant and animal tissues: Weigh about 0.1 g of sample, add 1 mL of buffer and fully grind. Centrifuge at 8000rpm and 4°C for 15 min. Dilute the supernatant with distilled water 10 times for test.
2. Fermentation broth: The fermentation broth is centrifuged at 8000 rpm, 4°C for 15 min, and the supernatant is taken as a sample to be tested.
3. Enzyme dry powder: Weigh about 1mg, add 1mL of buffer to dissolve, Centrifuge at 8000rpm and 4°C for 15 min, and dilute with distilled water 10 times for testing.

Procedure:

1. Preheat spectrophotometer for 30min, adjust the wavelength to 540 nm and set the counter to zero with distilled water.
2. Add reagent to a 1.5mL EP tube:

Reagent name (μL)	Control tube (Ac)	Test tube (At)	Blank tube (Ab)	Standard tube (As)
Sample	200	200		
2μmol/mL Standard				200
Distilled water			200	
buffer	300	300	300	300
Reagent 1		200	200	200
Mix well, react in a water bath at 50°C for 30 min, and inactivate immediately in a boiling water bath for 10 min. (Be careful not to let the lid pop open, so as not to enter the water and change the reaction system)				
Reagent 1	200			
Reagent 2	300	300	300	300

Mix well, develop color for 5 min in a boiling water bath (be careful not to open the lid to prevent water from changing the reaction system). After cooling, absorb 200μL in a 96-well plate or cuvette, and measure the absorbance at 540nm of each tube as soon as possible. Recorded as A measurement, A_C , A_S , A_B . Calculate $\Delta A = A_t - A_c$, $\Delta A_s = A_s - A_b$.

Calculation:

1. Fermentation broth:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme that catalyzes the production of 1μmol reducing sugar per minutes every mL fermentation broth in the reaction system.

$$BAX (U/mg mL) = C_s \times \Delta A \div \Delta A_s \div T = 0.067 \times \Delta A \div \Delta A_s$$

C_s : standard concentration , 2μmol/mL;

T : reaction time, 30min;

2. Enzyme dry powder:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme that catalyzes the production of 1μmol reducing sugar per minutes every mg powder in the reaction system.

$$BAX (U/mg prot) = 10 \times C_s \times \Delta A \div \Delta A_s \times V_e \div W_1 \div T = 0.67 \times \Delta A \div \Delta A_s \div W_1$$

10 : Sample dilution factor;

C_s : standard concentration , 2μmol/mL;

T : reaction time, 30min;

V_e : buffer volume , 1mL;

W_1 : enzyme dry powder weight , mg;

3. Tissue:

(1) Sample weight:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme that catalyzes the production of 1 μ mol reducing sugar per minute every gram tissue in the reaction system.

$$\text{BAX (U/g)} = 10 \times C_s \times \Delta A \div \Delta A_s \times V_e \div W_2 \div T = 0.67 \times \Delta A \div \Delta A_s \div W_2$$

(2) Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme that catalyzes the production of 1 μ mol reducing sugar per min every mg protein in the reaction system.

$$\text{BAX (U/mg prot)} = 10 \times C_s \times \Delta A \div \Delta A_s \times V_s \div (V_s \times C_{pr}) \div T = 0.67 \times \Delta A \div \Delta A_s \div C_{pr}$$

10 : Sample dilution factor;

C_s : standard concentration , 2 μ mol/mL;

T : reaction time, 30min;

V_e : buffer volume , 1mL;

W₂ : sample weight , g;

C_{pr} : protein concentration , mg/mL;

V_s : sample volume , 0.2mL.

Precautions:

The change in absorbance should be controlled between 0.01 and 1, otherwise increase the sample volume or dilute the sample. Note that the dilution factor involved in the calculation should be changed accordingly.

Experimental examples:

1. Take 0.1 g of leaves and add 1 mL of Buffer for sample processing. Take the supernatant and dilute ten times with distilled water and follow the measurement procedure. Calculate $\Delta A = A_t - A_c = 0.924 - 0.671 = 0.253$, $\Delta A_s = A_s - A_b = 0.761 - 0.163 = 0.598$. The enzyme activity is calculated according to the sample mass.

$$\text{BAX (U/g)} = 0.67 \times \Delta A \div \Delta A_s \div W_2 = 2.835 \text{ U/g.}$$

2. Take the yeast by centrifugation and take the supernatant and follow the determination procedure. Calculate $\Delta A = A_t - A_c = 1.135 - 1.001 = 0.134$, $\Delta A_s = A_s - A_b = 0.761 - 0.163 = 0.598$. The enzyme activity is calculated according to the sample mass.

$$\text{BAX (U/mL)} = 0.067 \times \Delta A \div \Delta A_s = 0.015 \text{ U/mL.}$$

Related products:

- AK0199/AK0198 Acidic Xylanase Activity Assay Kit
AK0201/AK0200 Neutral Xylanase Activity Assay Kit
AK0197/AK0196 β -xylosidase Activity Assay Kit