

## Soil nitrite reductase (S-NiR) Assay Kit

**Note:** Take two or three different samples for prediction before test.

**Operation Equipment:** Spectrophotometer/ Microplate reader

**Catalog Number:** AK0162

**Size:**100T/48S

### Components:

Reagent 1: Powder×1, storage at 4°C . Dissolve with 1mL of distilled water before use. The reagent can be saved for 2 weeks at 4°C . Dilute 400 times with distilled water before use, prepared when the solution will be used.

Reagent 2: Powder×1, storage at 4°C . Dissolve with 15mL of distilled water before use. The reagent can be saved for 2weeks at 4°C .

Reagent 3:15 mL×1, storage at 4°C . This solution is a saturated solution, just use the supernatant

Reagent 4:15 mL×1, storage at RT and protected from light.

Reagent 5:15 mL×1, storage at RT and protected from light.

Standard: 1 mL×1, storage at 4°C . 10 μmol/mL ofNaNO<sub>2</sub> standard solution.

### Product Description:

Soil nitrite reductase (S-NiR) is one of the key enzymes in denitrification. It is a reductase produced by soil denitrifying bacteria. It can reduce NO<sub>2</sub><sup>-</sup> to NO. The activity reflects the conversion efficiency of nitrogen in the process of biodegradation, and provides a certain basis for the study of nitrogen conversion.

Nitrite reductase can reduce NO<sub>2</sub><sup>-</sup> to NO, and reduce the NO<sub>2</sub><sup>-</sup> in the sample to participate in the diazotization reaction to produce a purple-red compound, that is, the change in absorbance at 540nm can reflect the activity of nitrite reductase in soil.

### Reagents and Equipment Required but Not Provided:

Spectrophotometer/ Microplate reader, adjustable transferpettor, balance, mortar/homogenizer, centrifuge, micro glass cuvette/ 96-well flat-bottom plate, sieve (30-50 mesh, or smaller), ice and distilled water.

### Procedure:

#### I . Sample preparation

Fresh soil samples are naturally air-dried or oven dried at 37°C and sieved through 30-50 mesh.

#### II . Determination

1. Preheat spectrophotometer/microplate reader for 30 min, adjust the wavelength to 540 nm and set the counter to zero with distilled water.
2. Dilute the standard solution with distilled water to prepare 0.8 、 0.6 、 0.4 、 0.2 、 0.1 、 0.05 μmol/mL standard solution.

3. Add reagent to a 1.5 mL EP tube:

	Non-matrix tube (An)	Blank tube1 (Ab1)	Control tube (Ac)	Test tube (At)	Standard tube (As)	Blank tube (Ab)
sample (g)	-	-	0.05	0.05	-	-
Distilled water (μL)	-	100	100	-	-	-
Reagent 1 (μL)	100	-	-	100	-	-
Reagent 2 (μL)	100	100	100	100	-	-
After mixing, react at 25°C for 3 h						
Reagent 3 (μL)	100	100	100	100	-	-
Fully shake for 30s, centrifuge at 10000 rpm for 10 min at 4°C .						
Supernatant (μL)	100	100	100	100	-	-
Standard (μL)	-	-	-	-	100	-
Reagent 4 (μL)	100	100	100	100	100	100
Reagent 5 (μL)	100	100	100	100	100	100
Distilled water (μL)						100

Mix well and react at room temperature for 15min. Take 200 μL into a micro glass cuvette/96 well plate and measure the absorbance value at the wavelength of 540nm, and record them as An, Ab1, Ac, At, As and Ab, and calculate  $\Delta A = (A_n - A_{b1}) - (A_t - A_c)$ ,  $\Delta A_s = A_s - A_b$ . Non-matrix tube (An), Blank tube1 (Ab1), Blank tube (Ab) only need to be done 1-2 times.

**III. Calculation:**

1. According to concentration of standard solution and absorbance to create the standard curve, take standard solution as X-axis,  $\Delta A_s$  as Y-axis. Take  $\Delta A$  into the equation to obtain x (μmol/mL)

**2. Fermentation broth:**

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the reduction of 1 μmol NO<sub>2</sub><sup>-</sup> per day every gram soil in the reaction system.

$$S\text{-NiR (U/g)} = x \times V_r \div T \div W = 2.4 \times x \div W.$$

T : reaction time, 3h=1/8 d;

V1: Enzymatic reaction volume, 0.3 mL;

W: soil weight, g;

**Related Products :**

AK0161/AK0160	Soil Hydroxylamine Reductase Activity Assay Kit
AK0329/AK0328	Soil Lignin peroxidase(S-Lip) Activity Assay Kit
AK0118/AK0117	Soil β- 1,4-Glucanase Activity Assay Kit
AK0120/AK0119	Soil Leucine Arylamidase (S-LAP) Activity Assay Kit
AK0574/AK0573	Soil Saccharase( S- SC) Activity Assay Kit



SunLong Biotech Co.,LTD  
Tel: 0086-571-56623320 Fax:0086-571-56623318  
E-mail:sales@sunlongbiotech.com  
www.sunlongbiotech.com

---

AK0370/AK0369 Soil Nitrate Reductase (NR) Activity Assay Kit