

Glutathione Peroxidase (GSH-Px/GPX) Activity Assay Kit

Note: Take two or three different samples for prediction before test.

Operation Equipment:Spectrophotometer

Cat No: AK0474

Size: 50T/24S

Components:

Extract solution: 30 mL×1, Storage at 4°C .

Reagent I: Powder×2, Storage at 4°C . Add 3.3 mL of Diluent to dissolve when the solution will be used.

Reagent II: 10 μL×1, Storage at 4°C . Add the liquid to the EP tube inside the reagent bottle. Dilute reagent II with the ratio of 2μL reagent II and 10 mL distilled water before use.

Reagent III: 60 mL×1, Storage at 4°C . If the bottom of the bottle is crystallized, it can be dissolved in water bath at 50C. This solution is a saturated solution. If the bottom of the bottle is still crystallized, the supernatant can be absorbed and used.

Reagent IV: 30 mL×1, Storage at 4°C .

Reagent V: 10mL×1, Storage at 4°C .

Standard: Powder×1, Storage at 4°C . 10 mg reduced glutathione (GSH). Add 0.405 mL of distilled water to the standard solution of 80 μmol/mL when the solution will be used.

Diluent: 20 mL×1, Storage at 4°C .

Product Description:

Glutathione peroxidase (glutathione peroxidase, GSH-Px or GPX) is an important peroxidase widely existed in the body. GPX can catalyzes the formation of oxidized glutathione (GSSG) from reduced glutathione (GSH) and reduce toxic hydrogen peroxide to non-toxic hydroxyl compounds.

GPX catalyzes the oxidation of GSH by hydrogen peroxide to produce GSSG. GSH can react with DTNB to form compounds with characteristic absorption peaks at 412 nm. The decrease of absorbance at 412 nm can reflect the activity of GPX.

Reagents and Equipment Required but Not Provided:

Spectrophotometer, balance, table centrifuge, 1 mL glass cuvette, mortar/homogenizer, EP tube.

Procedure

I. Sample preparation:

1. Tissue:

Accordance ratio Tissue weight (g): Extract solution (mL)=1:5~10 (Suggested 0.05g of tissue with 1mL of Extract solution), homogenate on ice bath. Centrifuge at 5000 rpm at 4°C for 10minutes, take the supernatant and place it on ice for test (If the supernatant is not clear, centrifuge for 3 minutes).

2. Bacteria or cells

Amount of cells (10^4): Extract solution (mL): 500~1000:1(Add 1mL of Extract solution to 5 million cells), ultrasonic with ice bath to break cells(300W,3s, interval 7s,total time 3minutes). Centrifuged at 5000 rpm at 4°C for 10minutes, take the supernatant and place it on ice for test (If the supernatant is not clear, centrifuge for 3 minutes).

3. Serum sample:

Detect directly.

II. Determination procedure:

1. Preheat spectrophotometer for 30minutes,adjust wavelength to 412 nm,set zero with distilled water.
2. The standard solution of 20 $\mu\text{mol/mL}$ was diluted to 0.25 $\mu\text{mol/mL}$ with the extraction solution. The standard solution of 100 μL is mixed with 400 μL of Reagent IV, and the concentration of the standard solution was 0.05 $\mu\text{mol/mL}$. The standard solution is prepared when the solution mixture will be used.
3. Mix the 150 μL of sample with the 150 μL of reagent I and place it at room temperature for 5 minutes.
4. Operation table: (1.5 mL centrifugal tube with the following reagents in turn).

Reagent Name(μL)	Test tube (T)	Control tube (C)
Sample Supernatant	100	-
Reagent I	100	100
Preheat for 5minutes at 37°C		
Reagent II	50	50
Reaction for 5 minutes at 37°C		
Reagent III	1000	1000
Sample mixtures	-	100

Centrifuge at 4000 rpm at room temperature for 5 minutes and take the supernatant into EP tube.

Reagent Name(μL)	Test tube (T)	Control tube (C)	Standard tube (S)	Black tube (B)
Diluent	-	-	-	500
Supernatant	500	500	-	-
Standard mixtures	-	-	500	-
Reagent IV	500	500	500	500
Reagent V	125	125	125	125

Well mix. then placed at room temperature for 15 minutes, the absorbance at 412 nm is measured. The absorbance is recorded as A_T , A_C , A_S and A_B , respectively. Calculate $\Delta A_T = A_C - A_T$, $\Delta A_S = A_S - A_B$.

III. Calculation:

1. Calculation of inhibition percentage

$$\text{Inhibitory percentage} = (A_C - A_T) / (A_C - A_B) \times 100\%$$

As far as possible, the inhibition percentage of the sample is within the range of 30-70%, and the closer it is to 50%, the more accurate it is. If inhibition percentage is less than 30% or more than 70%, it is usually necessary to adjust the dosage and re-determine it. If inhibition percentage is high, the sample should be diluted properly. If inhibition percentage is low, the sample with high concentration should be prepared again.

2. Calculation of GPX activity

1) Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the oxidation of 1 nmol of GSH per minute in the reaction system every milligram of protein.

$$\text{GPX (U/mg prot)} = \frac{\Delta A_T}{(\Delta A_S - C_S)} \times 1000 \times \frac{V_{EV}}{(C_{pr} \times V_{SV})} \div T = 200 \times \frac{\Delta A_T}{\Delta A_S - C_{pr}}$$

2) Sample weight

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the oxidation of 1 nmol of GSH per minute in the reaction system every gram of sample.

$$\text{GPX (U/g weight)} = \frac{\Delta A_T}{(\Delta A_S - C_S)} \times 1000 \times \frac{V_{EV}}{(V_{SV} \div V_{TV} \times W)} \div T = 200 \times \frac{\Delta A_T}{\Delta A_S - C_S} \div W$$

3) Cell amount

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the oxidation of 1 nmol of GSH per minute in the reaction system every 10^4 cells.

$$\text{GPX (U/10}^4\text{cell)} = \frac{\Delta A_T}{(\Delta A_S - C_S)} \times 1000 \times \frac{V_{EV}}{(N \times V_{SV} \div V_{TV})} \div T = 200 \times \frac{\Delta A_T}{\Delta A_S - C_S} \div N$$

4) Liquid volume:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the oxidation of 1 nmol of GSH per minute in the reaction system every milliliter of liquid.

$$\text{GPX (U/mL)} = \frac{\Delta A_T}{(\Delta A_S - C_S)} \times 1000 \times \frac{V_{EV}}{V_S} \div T = 200 \times \frac{\Delta A_T}{\Delta A_S - C_S}$$

C_S : Concentration of standard mixtures, 0.08 $\mu\text{mol/mL}$;

V_{EV} : Volume of enzymatic reaction system, 1.25 mL;

V_{sv} : Sample volume contained in sample mixtures, 0.1 mL;

V_{TV} : Extraction solution volume, 1 mL;

C_{pr} : Supernatant protein concentration, mg/mL;

T : Reaction time, 5 minutes;

N : The amount of cells, tens of thousands;

W : Sample weight, g;

1000: 1 μmol =1000 nmol.

Note:

1. When the absorbance is greater than 1.2, it is suggested that the sample be determined after diluted with the extraction solution.

2. It is recommended that not to take too many samples at a time, to avoid the influence of too long testing time on color development, which may let the determination is not accurate.

Experimental instances:

1. Take 0.1 g of mouse liver, add 1 mL of extract solution, homogenate and grind. Take the supernatant, dilute it by 40 times and test according to the measured steps. Calculate $A_T=0.108$, $A_C=0.303$, $A_S=0.491$, $A_B=0.033$, $\Delta A_T=A_C-A_T=0.195$, $\Delta A_S=A_S-A_B=0.458$, calculate the enzyme activity according to sample weight:

$$\text{GPX (U/g weight)} = 200 \times \frac{\Delta A_T}{\Delta A_S} \div W \times 40 \text{ (dilution ratio)} = 34061 \text{ U/g.}$$

2. Take 0.1 g of poplar leaf, add 1 mL of extract solution, homogenate and grind. Calculate $A_T=0.220$,

$A_C=0.318$, $A_S=0.491$, $A_B=0.033$, $\Delta A_T=A_C-A_T=0.098$, $\Delta A_S=A_B-A_B=0.458$, calculate the enzyme activity according to sample weight:

$$\text{GPX (U/g weight)}=200 \times \Delta A_T \div \Delta A_S \div W=428 \text{ U/g.}$$

Recent Product citations

- [1] Yang Yang, Li Jing, Wei Cong, et al. Amelioration of nonalcoholic fatty liver disease by swertiamarin in fructose-fed mice. *Phytomedicine*. June 2019; 59.(IF4. 18)
- [2] Xuejuan Xia, Yuxiao, Xing, Guannan Li, et al. Antioxidant activity of whole grain Qingke (Tibetan *Hordeum vulgare* L) toward oxidative stress in d-galactose induced mouse model. *Journal of Functional Foods*. June 2018;(IF3. 197)
- [3] Qilong Wang, Guosheng Xiao, Guoliang Chen, et al. Toxic effect of microcystin-LR on blood vessel development. *Toxicological & Environmental Chemistry*. Feb 2019;(IF3.547)
- [4] Wang H, Li Y Y, Qiu L Y, et al. Involvement of DJ 1 in ischemic preconditioning induced delayed cardioprotection in vivo[J]. *Molecular medicine reports*, 2017, 15(2): 995- 1001

Related products:

AK0478/ AK0477	Reduced Glutathione (GSH) Assay Kit
AK0476/AK0475	Oxidized Glutathione (GSSG) Assay Kit
AK0482/AK0481	Oxidized Thioredoxin Reductase (TrxR) Assay Kit
AK0472/AK0471	γ -glutamate-cysteine ligase (GCL) Assay Kit