

Catalase (CAT) Activity Assay Kit

Note: It is necessary to predict 2-3 large difference samples before the formal determination.

Operation Equipment: Spectrophotometer/ Microplate reader

Cat No: AK0579

Size: 100T/96S

Components:

Extraction reagent: Liquid 110 mL×1. Storage at 4°C;

Reagent I: Liquid 30 mL×1. Storage at 4°C .

Reagent II : Liquid 110 μL×1. Storage at 4°C . Centrifuge before use.

CAT working reagent: Add 25 μL of Reagent II to 5 mL of Reagent I before use, mix thoroughly as Working solution (about 26T). Or according to the proportion of preparation, the reagent should be prepared just before use.

Product Description:

CAT is an enzyme found broadly in animals, plants, microorganisms and cultured cells. It is the main enzyme of clearing H₂O₂, which plays an important role in the active oxygen scavenging system.

H₂O₂ has characteristic absorption peak at 240 nm. It can be decomposed into water and oxygen by CAT which makes the absorbance of reagent at 240 nm decreases. The activity of CAT can be calculated according to the change rate of absorbance.

Reagents and Equipment Required but Not Provided:

Spectrophotometer/microplate reader, refrigerated centrifuge, transferpeltor, micro quartz cuvette/96 well UV flat-bottom plate, mortar/ homogenizer, ice and distilled water.

Procedure:

I. Sample preparation:

1. Bacteria or cells:

Collect bacteria or cells into the centrifuge tube, after centrifugation discard supernatant. It is suggested that add 1 mL of Extraction reagent to 5 million of bacteria or cells. Use ultrasonication to split bacteria and cells (place on ice, ultrasonic power 200W, working time 3 seconds, interval 10 seconds, repeat for 30 times). Centrifuge at 8000 ×g for 10 minutes at 4°C to remove insoluble materials and take the supernatant on ice for testing.

2. Tissue:

It is suggested that add 1 mL of Extraction reagent to 0.1 g of tissue, and fully homogenize on ice bath. Centrifuge at 8000 ×g for 10 minutes at 4°C to remove insoluble materials, and take the supernatant on ice for test.

3. Serum (plasma) sample: Detect sample directly.

II. Determination procedure:

1. Preheat the spectrophotometer more than 30 minutes, adjust the wavelength to 240 nm, set zero with distilled water.
2. Preheat CAT working reagent in water bath at 37°C(mammals) or 25°C (other species) for 10 minutes.
3. Add 190 μ L of CAT working reagent and 10 μ L of sample in micro quartz cuvette/96 well UV flat-bottom plate. Immediately mix and detect the absorbance at 240 nm at the initial time(A1) and the absorbance after reaction for 1 minute(A2), calculate $\Delta A=A1-A2$.

III. Calculation:

A. micro quartz cuvette

1. Serum (plasma) sample

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the degradation of 1 μ mol ofH₂O₂ in the reaction system per minute every milliliter serum (plasma).

$$\text{CAT (U/mL)} = [\Delta A \times V_{rv} \div (\epsilon \times d) \times 10^6] \div V_s \div T = 459 \times \Delta A$$

2. Tissue, bacteria or cells

1) Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the degradation of 1 μ mol ofH₂O₂ in the reaction system per minute every milligram protein.

$$\text{CAT (U/mg prot)} = [\Delta A \times V_{rv} \div (\epsilon \times d) \times 10^6] \div (V_s \times C_{pr}) \div T = 459 \times \Delta A \div C_{pr}$$

2) Sample weight

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the degradation of 1 μ mol ofH₂O₂ in the reaction system per minute every gram tissue sample.

$$\text{CAT (U/g weight)} = [\Delta A \times V_{rv} \div (\epsilon \times d) \times 10^6] \div (W \times V_s \div V_{sv}) \div T = 459 \times \Delta A \div W$$

3) Bacteria or cells

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the degradation of 1 μ mol ofH₂O₂ in the reaction system per minute every 10⁴ bacteria or cells.

$$\text{CAT (U/10}^4\text{cell)} = [\Delta A \times V_{rv} \div (\epsilon \times d) \times 10^6] \div (500 \times V_s \div V_{sv}) \div T = 0.917 \times \Delta A$$

V_{rv}: Reaction total volume, 2 \times 10⁻⁴ L;

ϵ : Molar extinction coefficient, 43.6 L/mol/cm;

d: Light path of cuvette, 1 cm;

V_s: Sample volume,0.01 mL;

V_{sv}: Extraction volume, 1 mL;

T: Reaction time, 1 minute;

C_{pr}: Sample protein concentration, mg/mL;

W: Sample weight, g;

500: Total number of bacteria and cells, 5 million;

10^6 : Unit conversion factor, 1 mol= 10^6 μ mol.

B. 96 well UV flat-bottom plate

1. Serum (plasma) sample

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the degradation of 1 μ mol of H_2O_2 in the reaction system per minute every milliliter serum (plasma).

$$CAT (U/mL) = [\Delta A \times V_{rv} \div (\epsilon \times d) \times 10^6] \div V_s \div T = 764.5 \times \Delta A$$

2. Tissue, bacteria or cells

1) Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the degradation of 1 μ mol of H_2O_2 in the reaction system per minute every milligram protein.

$$CAT (U/mg prot) = [\Delta A \times V_{rv} \div (\epsilon \times d) \times 10^6] \div (V_s \times C_{pr}) \div T = 764.5 \times \Delta A \div C_{pr}$$

2) Sample weight

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the degradation of 1 μ mol of H_2O_2 in the reaction system per minute every gram tissue sample.

$$CAT (U/g) = [\Delta A \times V_{rv} \div (\epsilon \times d) \times 10^6] \div (W \times V_s \div V_{sv}) \div T = 764.5 \times \Delta A \div W$$

3) Bacteria or cells

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the degradation of 1 μ mol of H_2O_2 in the reaction system per minute every 10^4 bacteria or cells.

$$CAT (U/10^4 cell) = [\Delta A \times V_{rv} \div (\epsilon \times d) \times 10^6] \div (500 \times V_s \div V_{sv}) \div T = 1.529 \times \Delta A$$

V_{rv} : Reaction total volume, 2×10^{-4} L;

ϵ : Molar extinction coefficient, 43.6 L/mol/cm;

d : light path of 96 well plate, 0.6 cm;

V_s : Sample volume, 0.01 mL;

V_{sv} : Extraction volume, 1 mL;

T : Reaction time, 1 minute;

C_{pr} : Sample protein concentration, mg/mL;

W : Sample weight, g;

500: Total number of bacteria and cells, 5 million;

10^6 : Unit conversion factor, 1 mol= 10^6 μ mol.

Note:

If there are a lot of bubbles in the reaction solution, dilute the sample with distilled water before determination.

Recent Product Citations:

[1] Zhang Z, Liu H, Sun C, et al. A C₂H₂ zinc-finger protein OsZFP213 interacts with OsMAPK₃ to enhance salt tolerance in rice[J]. Journal of plant physiology, 2018, 229: 100- 110.

[2] Yin Y J, Chen C J, Guo S W, et al. The fight against Panax notoginseng root-rot disease using zingiberaceae essential oils as potential weapons[J]. Frontiers in plant science, 2018, 9: 1346.

[3] Yang Y, Li J, Wei C, et al. Amelioration of nonalcoholic fatty liver disease by swertiamarin in fructose-fed mice[J]. Phytomedicine, 2019, 59: 152782.

[4] Chen G, Jia Z, Wang L, et al. Effect of acute exposure of saxitoxin on development of zebrafish embryos (Danio rerio) [J]. Environmental Research, 2020: 109432.

References:

[1] Catalase in vitro. [J]. Methods Enzymol, 105:121- 126.

[2] Johansson L H, Borg L A H. A spectrophotometric method for determination of catalase activity in small tissue samples[J]. Analytical biochemistry, 1988, 174(1): 331-336.

Related Products:

AK0293/AK0292 Polyphenol Oxidase(PPO) Activity Assay Kit

AK0578/AK0577 Phenylalnine Ammonialyase(PAL) Activity Assay Kit

AK0584/AK0583 Superoxide Dismutase(SOD) Activity Assay Kit

AK0598/AK0597 Peroxidase(POD) Activity Assay Kit