

Active Insulin Like Growth Factor Binding Protein 3 (IGFBP3) Instruction Manual

SBPA034Hu01

Homo sapiens (Human)

Buffer Formulation	PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	9.3
Applications	Cell culture; Activity Assays.

ACTIVITY TEST




Figure 1. The binding activity of IGFBP3 with FN

Insulin-like growth factor-binding protein 3, also known as IGFBP3 is one of six IGF binding proteins (IGFBP1 to IGFBP6) that have highly conserved structures and bind the insulin-like growth factors IGF-1 and IGF-2 with high affinity. Within tissues, IGFBP3

can bind IGF1 and IGF2 released by many cell types, and block their access to the IGF-1 receptor (IGF1R), which is activated by both IGFs. IGFBP3 also interacts with cell-surface proteins, affecting cell signaling from outside the cell or after internalization, and also enters the cell nucleus where it binds to nuclear hormone receptors and other ligands. Besides, Fibronectin (FN) has been identified as an interactor of IGFBP3, thus a binding ELISA assay was conducted to detect the interaction of recombinant human IGFBP3 and recombinant human FN. Briefly, IGFBP3 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100ul were then transferred to FN-coated microtiter wells and incubated for 1h at 37°C. Wells were washed with PBST and incubated for 1h with anti-IGFBP3 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of IGFBP3 and FN was shown in Figure 1, the EC50 was 78 ng/ml.

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE




Figure. Western Blot

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.