

Active Macrophage Inflammatory Protein 3 Alpha (MIP3 α) Instruction Manual

SBPA064Hu01

Homo sapiens (Human)

Buffer Formulation	PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	9.6
Applications	Cell culture; Activity Assays.

ACTIVITY TEST




Figure 1. The chemotactic effect of recombinant human MIP-3 α on Jurkat cells

(A) Jurkat cells were seeded into the upper chambers and serum free RPMI 1640 with 10ng/mL MIP-3 α was added in lower chamber, then cells in lower chamber were observed at low magnification ($\times 100$) after incubation for 2h;

(B) Jurkat cells were seeded into the upper chambers and serum free RPMI 1640 without MIP-3 α was added in lower chamber, then cells in lower chamber were observed at low magnification ($\times 100$) after incubation for 2h.

Macrophage Inflammatory Protein 3 Alpha (MIP-3 α), also known as LARC (liver and activation-regulated chemokine), Exodus-1 or CCL20, is a CC chemokine with a selective chemotactic activity for lymphocytes and dendritic cells (DCs). MIP3 α is

produced by activated cells, including monocytes, T cells, endothelial cells, epithelial cells, and fibroblasts and is expressed in liver, lung, and some lymphoid tissues. This chemokine elicits its effects on its target cells by binding to the chemokine receptor CCR7. It attracts certain cells of the immune system, including dendritic cells and antigen-engaged B cells, CCR7 central-memory T-Cells. Thus, chemotaxis assay used 24-well microchemotaxis system was undertaken to detect the chemotactic effect of recombinant human MIP-3 α on the Jurkat cell line. Briefly, Jurkat cells were seeded into the upper chambers (200 μ L cell suspension, 10⁶ cells/mL in RPMI 1640 with FBS free) and MIP-3 α (0.01ng/mL, 0.1ng/mL, 1ng/mL, 10ng/mL, 100ng/mL and 1000ng/mL diluted separately in serum free RPMI 1640) was added in lower chamber with a polycarbonate filter (8 μ m pore size) used to separate the two compartments. After incubation at 37°C with 5% CO₂ for 2h, the filter was removed, then cells in low chamber were observed by inverted microscope at low magnification (\times 100) and the number of migrated cells were counted at high magnification (\times 400) randomly (five fields for each filter). Result shows MIP-3 α is able to induce migration of Jurkat cells. The migrated Jurkat cells in low chamber at low magnification (\times 100) were shown in Figure 1. Five fields of each chamber were randomly chosen, and the migrated cells were counted at high magnification (\times 400). Statistical results were shown in Figure 2. The optimum chemotaxis of recombinant human MIP-3 α occurs at 1-10ng/mL.




Figure 2. The chemotactic effect of recombinant human MIP-3 α on Jurkat cells

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. Gene Sequencing (Extract)

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.