

Active Stromal Cell Derived Factor 1 (SDF1) Instruction Manual

SBPA077Hu02

Homo sapiens (Human)

Buffer Formulation

20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,
1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits

Freeze-dried powder

Purity

> 97%

Isoelectric Point

8.5

Applications

Cell culture; Activity Assays.

ACTIVITY TEST




Figure 1. The chemotactic effect of SDF1 on THP1 cells.

(A) THP-1 cells were seeded into the upper chambers and serum free RPMI 1640 with 10ng/mL SDF1 was added in lower chamber, then cells in lower chamber were observed at low magnification ($\times 100$) after incubation for 3h;

(B) THP-1 cells were seeded into the upper chambers and serum free RPMI 1640 without SDF1 was added in lower chamber, then cells in lower chamber were observed at low magnification ($\times 100$) after incubation for 3h.




Figure 2. The chemotactic effect of SDF1 on THP-1 cells

SDF1 (stromal cell-derived factor 1), also known as C-X-C motif chemokine 12, is a chemokine protein that has chemotaxis active on T-lymphocytes and monocytes. It is thought that SDF1 stimulates migration of monocytes and T-lymphocytes through its receptors, CXCR4 and ACKR3; thus, chemotaxis assay used 24-well microchemotaxis system was undertaken to detect the chemotactic effect of SDF1 on the human monocytic cell line THP-1. Briefly, THP-1 cells were seeded into the upper chambers (100uL cell suspension, 106 cells/mL in RPMI 1640 with 0.5% FBS) and SDF1 (10ng/mL, 30ng/mL and 60ng/mL diluted separately in serum free RPMI 1640) was added in lower chamber with a polycarbonate filter (8µm pore size) used to separate the two compartments. After incubation at 37°C with 5% CO₂ for 3h, the filter was removed, then cells in low chamber were observed by inverted microscope at low magnification (×100) and the number of migrated cells were counted at high magnification (×400) randomly (five fields for each filter). Result shows SDF1 is able to induce migration of THP-1 cells. The migrated THP-1 cells in low chamber at low magnification (×100) were shown in Figure 1. Five fields of each chamber were randomly chosen, and the migrated cells were counted at high magnification (×400). Statistical results were shown in Figure 2. The optimum chemotaxis of SDF1 occurs at 10ng/mL.

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image




Figure. Western Blot; Sample: Recombinant SDF1, Human.

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.