Active Interleukin 1 Receptor Antagonist (IL1RA) Instruction Manual

SBPA103Hu01

Homo sapiens (Human)

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 95% Isoelectric Point 6.0

Applications Cell culture; Activity Assays.

ACTIVITY TEST




Figure 1. The binding activity of IL1RA with IL1R1.

IL1RA (interleukin-1 receptor antagonist) is an agent that binds to the cell surface interleukin-1 receptor (IL-1R), which would prevent IL-1 from intracellular signal transduction. Thus, a binding ELISA assay was constructed to detect the association of recombinant human IL1RA with recombinant human IL1R1. Briefly, IL1RA were diluted serially in PBS with 0.1% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to IL1R1-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-IL1RA pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody,

wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add $50\mu L$ stop solution to the wells and read at 450nm immediately. The binding activity of IL1RA with IL1R1 was shown in Figure 1 and this effect was in a dose dependent manner.

USAGE


Reconstitute in 20mM Tris, 150mM NaCl (PH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.






Figure. Western Blot; Sample: Recombinant IL1RA, Human.

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.