

Active Interleukin 1 Beta (IL1b) Instruction Manual

SBPA147Mu01

Mus musculus (Mouse)

Buffer Formulation	PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	6.7
Applications	Cell culture; Activity Assays.

ACTIVITY TEST




Figure1. IL-8 levels in the cell supernatant of A549 induced by IL-1β

Interleukin 1 beta (IL-1β) also known as leukocytic pyrogen, leukocytic endogenous mediator, mononuclear cell factor, lymphocyte activating factor and other names, is a member of the interleukin 1 family of cytokines. This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. It has been reported that IL-1β can induced IL-8 production in A549 cells. To test the bioactivity of IL-1β, A549 cells were seeded into 24-well plate at a density of 1x10⁵ cells/mL, and allowed to attach overnight before treated with certain concentrations of IL-1β for 24h and IL-8 levels in the cell

supernatant were determined by ELISA. IL-8 levels in the cell supernatant of A549 cells increased significantly after stimulated with IL-1 β have shown in Figure1.

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.