

# Active Bone Morphogenetic Protein 7 (BMP7) Instruction Manual

## SBPA181Mu01

### Mus musculus (Mouse)

#### Buffer Formulation

20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

#### Traits

Freeze-dried powder

#### Purity

> 97%

#### Isoelectric Point

8.5

#### Applications

Cell culture; Activity Assays.

#### ACTIVITY TEST





Figure 1. Cell proliferation of MDA-MB-231 cells after stimulated with BMP7.

(A) MCF-7 cells cultured in DMEM, stimulated with 10ng/mL BMP7 for 72h;

(B) Unstimulated MDA-MB-231 cells cultured in DMEM for 72h.

BMP7 (Bone morphogenetic protein 7), which belongs to the TGF- $\beta$  superfamily, is a signaling molecule with the ability to promote bone formation. BMP7 has also been implicated in various types of cancer, including breast cancer. It has been reported that BMP7 treatment induced cell growth promotion of MDA-MB-231 breast cancer line. To test the effect of BMP7 on cell proliferation, MDA-MB-231 cells were seeded into triplicate wells of 96-well plates at a density of 2,000 cells/well and allowed to attach overnight, then the medium was replaced with serum-free standard DMEM prior to the addition of various concentrations of BMP7. After incubated for 72h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 $\mu$ L of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37°C.



**Figure 2. The dose-effect curve of BMP7 on MDA-MB-231 cells**

The dose-effect curve of BMP7 was shown in Figure 2. It was obvious that BMP7 significantly promoted cell proliferation of MDA-MB-231 cells. The ED50 for this effect is typically 3.483 to 9.017 ng/mL.

## USAGE

Reconstitute in 20mM Tris, 150mM NaCl (PH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## Image



SDS-PAGE Image




Figure. Western Blot; Sample: Recombinant BMP7, Mouse.

**[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.