

Active Fibroblast Growth Factor 11 (FGF11) Instruction Manual

SBPC320Hu01

Homo sapiens (Human)

Buffer Formulation

20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits

Freeze-dried powder

Purity

> 90%

Isoelectric Point

10.0

Applications

Cell culture; Activity Assays.

ACTIVITY TEST




Figure 1. The binding activity of FGF11 with FGFR1

Fibroblast Growth Factor 11 (FGF11) is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. The function of this gene has not yet been determined. Besides, Fibroblast Growth Factor Receptor 1 (FGFR1) has been identified as an interactor of FGF11, thus a binding ELISA assay was conducted to detect the interaction of recombinant human FGF11 and recombinant human FGFR1. Briefly, FGF11 were diluted serially in PBS, with 0.01% BSA (pH 7.4).

Duplicate samples of 100µl then transferred to FGFR1-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-FGF11 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µl stop solution to the wells and read at 450nm immediately. The binding activity of FGF11 and FGFR1 was shown in Figure 1, and this effect was in a dose dependent manner.

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (PH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.