

Active Interleukin 17 Receptor D (IL17RD) Instruction Manual

SBPD344Hu01

Homo sapiens (Human)

Buffer Formulation

20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits

Freeze-dried powder

Purity

> 90%

Isoelectric Point

8.9

Applications

Cell culture; Activity Assays.

ACTIVITY TEST




Figure 1. The binding activity of IL17RD with IL17RA.




Figure 2. The binding activity of IL17RD with IL17.

IL17RD (interleukin-17 receptor D) is a membrane protein belonging to the IL17R protein family, acting as a component of the interleukin-17 receptor signaling complex. Knowing that the interaction between this protein and IL-17R does not require the interleukin, we have conducted a binding ELISA assay to detect the interaction of recombinant human IL17RD with both recombinant human IL17RA and IL17. Briefly, IL17RD were diluted serially in PBS, with 0.01%BSA (pH 7.4). Duplicate samples of 100uL were then transferred to IL17RA-coated or IL17-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-IL17RA pAb and anti-IL17 pAb separately, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of IL17RD with IL17RA and IL17 was shown in Figure 1 and Figure 2 respectively and this effect was in a dose dependent manner.

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



SDS-PAGE Image




Figure. Western Blot; Sample: Recombinant IL17RD, Human.

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.