

Active Interleukin 17 Receptor C (IL17RC) Instruction Manual

SBPD345Hu01

Homo sapiens (Human)

Buffer Formulation

PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Traits

Freeze-dried powder

Purity

> 97%

Isoelectric Point

5.8

Applications

Cell culture; Activity Assays.

ACTIVITY TEST




Figure 1. The binding activity of IL17RC with IL17RA.

IL17RC (Interleukin-17 receptor C) is a receptor for IL17A and IL17F homodimers as part of a heterodimeric complex with IL17RA. This protein is expressed in nonhemopoietic tissues, and binds both IL-17A and IL-17F with similar affinities. A binding ELISA assay was conducted to detect the association of IL17RC with IL17RA. Briefly, IL17RC were diluted serially in PBS with 0.01%BSA (pH 7.4). Duplicate samples of 100uL were then transferred to IL17RA-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-IL17RC pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate

solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of IL17RC with IL17RA was shown in Figure 1 and this effect was in a dose dependent manner.

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE; Sample: Active recombinant IL17RC, Human.




Figure. Western Blot; Sample: Recombinant IL17RC, Human.

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

