

# Active Adenosylhomocysteinase (AHCY) Instruction Manual

## SBPG364Hu01

**Homo sapiens (Human)**

**Buffer Formulation**

20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Traits**

Freeze-dried powder

**Purity**

> 95%

**Isoelectric Point**

5.9

**Applications**

Cell culture; Activity Assays.

### ACTIVITY TEST




Figure. The binding activity of AHCY with MHCB.

Adenosylhomocysteinase (S-adenosylhomocysteine synthase, S-adenosylhomocysteine hydrolase, adenosylhomocysteine hydrolase, S-adenosylhomocysteinase, SAHase, AdoHcyase, AHCY) is an enzyme that converts S-adenosylhomocysteine to homocysteine and adenosine. The enzyme contains one tightly bound NAD per subunit. Besides, Major Histocompatibility Complex Class I B (MHCBI) has been identified as an interactor of AHCY, thus a binding ELISA assay was conducted to detect the interaction of recombinant human AHCY and recombinant human MHCBI. Briefly, AHCY were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to MHCBI-coated microtiter wells and incubated for 2h at 37°C. Wells were

washed with PBST and incubated for 1h with anti-AHCY pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of AHCY and MHCB was shown in Figure 1, and this effect was in a dose dependent manner.

## USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## Image




Figure. SDS-PAGE




Figure. Western Blot

**[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.