Eukaryotic Brain Derived Neurotrophic Factor (BDNF) Instruction Manual

SFPA007Hu62

Homo sapiens (Human)

Source Eukaryotic expression

Host 293F cell

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass27.4kDa

Accurate Molecular Mass 35&15kDa(Analysis of differences refer to the manual)

Residues & Tags Ala19~Arg247 with N-terminal His Tag **Buffer Formulation** PBS, pH7.4, containing 5% Trehalose.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 8.9

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

AP MKEANIRGQG GLAYPGVRTH GTLESVNGPK
AGSRGLTSLA DTFEHVIEEL LDEDQKVRPN EENNKDADLY TSRVMLSSQV
PLEPPLLFLL EEYKNYLDAA NMSMRVRRHS DPARRGELSV CDSISEWVTA
ADKKTAVDMS GGTVTVLEKV PVSKGQLKQY FYETKCNPMG YTKEGCRGID
KRHWNSQCRT TQSYVRALTM DSKKRIGWRF IRIDTSCVCT LTIKRGR

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.