

# Eukaryotic Interleukin 6 (IL6)

## Instruction Manual

**SFPA051Po61**

**Sus scrofa; Porcine (Pig)**

<b>Source</b>	Eukaryotic expression
<b>Host</b>	293F cell
<b>Endotoxin Level</b>	<1.0EU per 1ug (determined by the LAL method)
<b>Subcellular Location</b>	Secreted
<b>Predicted Molecular Mass</b>	22.5kDa
<b>Accurate Molecular Mass</b>	25/30kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Gly30~Met212 with N-terminal His Tag
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 97%
<b>Isoelectric Point</b>	5.8
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

### SEQUENCE

```
                E RLEEDAKGDA TSDKMLFTSP
DKTEELIKYI LGKISAMRKE MCEKYEKCEN SKEVLAENNL NLPKMAEKDG
CFQSGFNQET CLMRITGLV EFQIYLDYLQ KEYESNKGNV EAVQISTKAL
IQTLRQKGKN PDKATTPNPT TNAGLLDKLQ SQNEWMKNTK IILILRSLED
FLQFSLRAIR IM
```

### USAGE

Reconstitute in PBS or others.


### STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### Image



SDS-PAGE Image

### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.