

# Recombinant Interleukin 18 (IL18)

## Instruction Manual

**SIPA043Po01**

**Sus scrofa; Porcine (Pig)**

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Secreted
<b>Predicted Molecular Mass</b>	21.7kDa
<b>Accurate Molecular Mass</b>	22kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Tyr36~Asn192 with N-terminal His Tag
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 95%
<b>Isoelectric Point</b>	5.4
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

### SEQUENCE

```
                                YFGKL EPKLSIIRNL
NDQVLFINQG HQAVFEDMPD SDCSDNAPQT VFIIYMYKDS LTRGLAVTIS
VQCKKMSTLS CKNKTLSEFKE MSPPDNIDDE GNDIIFQRS VPGHDDKIQF
ESSLYKGYFL ACKKENDLFK LILKEKDECG DKSIMFTVQN KN
```

### USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.  
Do not vortex.

### STORAGE


Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## Image

TACTTTGGCAGCTTGAGCTTAACCTCATCATAGAACTCGAAGCAGTCTTTTCATTAACAGGACATCAGGCGGTGGAGGATATCCGATTCCTACTGTCAGATATGCACTCAGACGTATTATATATATATGATATAGATAGCCTCCTAG  
Y F G K L E P K L S I I R N L N D Q V L F I N Q G H Q A V F E D M P D S D C S D N A P Q T V F I I Y M Y K D S L T R



SDS-PAGE Image

## **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.