

Recombinant Tumor Necrosis Factor Alpha (TNFa) Instruction Manual

SIPA083Eq01

Equus caballus; Equine (Horse)

| | |
|---------------------------------|---|
| Source | Prokaryotic expression |
| Host | E.coli |
| Endotoxin Level | <1.0EU per 1µg (determined by the LAL method) |
| Subcellular Location | Membrane, Secreted |
| Predicted Molecular Mass | 20.9kDa |
| Accurate Molecular Mass | 21kDa(Analysis of differences refer to the manual) |
| Residues & Tags | Leu78~Leu234 with N-terminal His Tag |
| Buffer Formulation | 100mMNaHCO ₃ , 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300. |
| Traits | Freeze-dried powder |
| Purity | > 95% |
| Isoelectric Point | 6.8 |
| Applications | Positive Control; Immunogen; SDS-PAGE; WB. |

SEQUENCE

```
                LRS SSRTPSDKPV AHVVANPQAE
GQLQWLSGRA NALLANGVKL TDNQLVVPLD GLYLIYSQVL FKGQGCPSTH
VLLTHTISRL AVSYPSKVNL LSAIKSPCHT ESPEQAEAKP WYEPIYLGGV
FQLEKGDQLS AEINQPNYLD FAESGQVYFG IIAL
```

USAGE


Reconstitute in 100mM NaHCO₃, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



SDS-PAGE Image

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.