

# Recombinant Thrombopoietin (TPO) Instruction Manual

## SIPA085Hu01

### Homo sapiens (Human)

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Secreted
<b>Predicted Molecular Mass</b>	65.0kDa
<b>Accurate Molecular Mass</b>	65kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Ser22~Gly349 with N-terminal His and GST Tag
<b>Buffer Formulation</b>	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 97%
<b>Isoelectric Point</b>	8.5
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

### SEQUENCE

```
SPAPPACDL RVLSKLLRDS HVLHSRLSQC  
PEVHPLPTPV LLPAVDFSLG EWKTQMEETK AQDILGAVTL LLEGVMAARG  
QLGPTCLSSL LGQLSGQVRL LLGALQSLLG TQGRTTAHKD PNAIFLSFQH  
LLRGKVRFLM LVGGSTLCVR RAPPTTAVPS RTSLVLTLINE LPNRTSGLLE  
TNFTASARTT GSGLLKWQQG FRAKIPGLLN QTSRSLDQIP GYLNRIHELL  
NGTRGLFPGP SRRTLGPDI SSGTSDTGSL PPNLQPGYSP SPTHPTGQY  
TLFPLPPTLP TPVVQLHPLL PDPSAPTPTP TSPLLNTSYT HSQNLSQEG
```

### USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL.  
Do not vortex.


### STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### **Image**



SDS-PAGE Image

### **[IMPORTANT NOTE]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.