Recombinant Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL) Instruction Manual

SIPA086Ra01

Rattus norvegicus (Rat)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationMembranePredicted Molecular Mass52.0kDa

Accurate Molecular Mass 50kDa(Analysis of differences refer to the manual)

Residues & Tags Pro115~Ala287 with N-terminal His and GST Tag

Buffer Formulation PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Traits Freeze-dried powder

Purity > 80% Isoelectric Point 7.3

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

PPLPRG RRPQRVAAHI TGITRRSNLA LIPISKDGKT LGQKIETWES SRRGHSFLNH VHLRNGELVI QEEGLYYIYS QTYYRFKEAK EASKTVSKDG GRIKQMVQYI YKYTSYPDPI LLMKSARNSC WSREAEYGLY SIYQGGLFEL KENDRIFVSV TNEHLMDLDH EASFFGA

USAGE

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.