Recombinant Anti-Mullerian Hormone (AMH) Instruction Manual

SIPA104Hu02

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass44.3kDa

Accurate Molecular Mass 48kDa(Analysis of differences refer to the manual)

Residues & Tags Thr 193~Arg 560 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01%skl, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 8.8

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

				TRYLVLAV
DRPAGAWRGS	GLALTLQPRG	EDSRLSTARL	QALLFGDDHR	CFTRMTPALL
LLPRSEPAPL	PAHGQLDTVP	FPPPRPSAEL	EESPPSADPF	LETLTRLVRA
LRVPPARASA	PRLALDPDAL	AGFPQGLVNL	SDPAALERLL	DGEEPLLLLL
RPTAATTGDP	APLHDPTSAP	WATALARRVA	AELQAAAAEL	RSLPGLPPAT
APLLARLLAL	CPGGPGGLGD	PLRALLLLKA	LQGLRVEWRG	RDPRGPGRAQ
RSAGATAADG	PCALRELSVD	LRAERSVLIP	ETYQANNCQG	VCGWPQSDRN
PRYGNHVVLL	LKMQVRGAAL	ARPPCCVPTA	YAGKLLISLS	EERISAHHVP
NMVATECGCR				

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.






Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.