

# Recombinant Tissue Plasminogen Activator (tPA) Instruction Manual

## SIPA138Po01

**Sus scrofa; Porcine (Pig)**

<b>Source</b>	Prokaryotic expression
<b>Host</b>	E.coli
<b>Endotoxin Level</b>	<1.0EU per 1µg (determined by the LAL method)
<b>Subcellular Location</b>	Secreted
<b>Predicted Molecular Mass</b>	88.3KDa
<b>Accurate Molecular Mass</b>	88kDa(Analysis of differences refer to the manual)
<b>Residues &amp; Tags</b>	Val39~Pro562 with N-terminal His and GST Tag
<b>Buffer Formulation</b>	100mMNaHCO <sub>3</sub> , 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
<b>Traits</b>	Freeze-dried powder
<b>Purity</b>	> 90%
<b>Isoelectric Point</b>	7.2
<b>Applications</b>	Positive Control; Immunogen; SDS-PAGE; WB.

## SEQUENCE

```
VT CRDEKTQMIY
QQHQSWLRPL LRGNRVEHCW CNDGQTQCHS VPKVSCSEPR CFNGGTCLQA
IYFSDFVCQC PVGFIGRQCE IDARATCYED QGITYRGTWS TTESGAECVN
WNTSGLASMP YNGRRPDAVK LGLGNHNYCR NPKDKSKPWC YIFKAEKYSY
DFCSTPACTK EKEECYTGKG LDYRGTRSLT MSGAFCLPWN SLVLMGKIYT
AWNSNAQTLG LGKHNYCRNP DGDTQPWCHV LKDHKLTWEY CDLPQCVTCG
LRQYKEPQFR IKGGLYADIT SHPWQAAIFV KNRRSPGERF LCGGILISSC
WVLSAAHCFQ ERFPPHHVRV VLGRTYRLVP GEEEQAFEVE KYIVHKEFDD
DTYDNDIALL QLKSDSLTCA QESDAVRTVC LPEANLQLPD WTECELSGYG
KHEASSPFYS ERLKEAHVRL YPSSRCTSKH LFNKTITNNM LCAGDTRSGG
DNANLHDACQ GDSGGPLVCM KGNHMTLVGV ISWGLGCGQK DVPGVYTKVT
NYLNNWIRDNT RP
```

## USAGE

Reconstitute in 100mM NaHCO<sub>3</sub>, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


## STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

## STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## Image



SDS-PAGE Image

## [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.