

Recombinant Tissue Plasminogen Activator (tPA) Instruction Manual

SIPA138Ra01

Rattus norvegicus (Rat)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Secreted, Extracellular matrix
Predicted Molecular Mass	23.6kDa
Accurate Molecular Mass	33kDa(Analysis of differences refer to the manual)
Residues & Tags	Gly311~Asn487B with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 90%
Isoelectric Point	6.2
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
GGLFTDITSH PWQAAIFVKN KRSPGERFLC GGVLISSCWV  
LSAAHCFVER FPPHHLKVVL GR TYRVVPG E EQTFEIEKY IVHKEFDDDT  
YNDIALQL RSDSSQCAQE SSSVGTACL DP DVQLPDWT ECELSGYGKH  
EASSPFFSDR LKEAHVRLYP SSRCTSQHLLF NKTITSN
```

USAGE

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

STORAGE


Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image

GGAGGACTTTCAGACATCACACACCTTGCCAGGCGCCATCTTGTCAAGAGAGGCTCCAGGAGAGAGATTCTGTGTGAGGGGCTGATCAGTTCTCTGTGGGTGCTATCTGCGCGCCATCTCTTGTAGAGAGTTCCACCCATCATCTTAAAGTGTCTTGGGCGAGCATAC
G G I F T D I T S H P V Q A A I F V K N X R S P G E R F L C G G V L I S S C W V L S A A H C F V E R F P P H E L K V V L G R T Y



F

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.