# Recombinant Cluster Of Differentiation 14 (CD14) Instruction Manual

# SIPA163Po01

## Sus scrofa; Porcine (Pig)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Membrane
Predicted Molecular Mass	41.3kDa
Accurate Molecular Mass	41kDa(Analysis of differences refer to the manual)
Residues & Tags	Thr21~Ala373 with N-terminal His Tag
Buffer Formulation	100mMNaHCO <sub>3</sub> , 500mMNaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Traits	Freeze-dried powder
Purity	> 97%
Isoelectric Point	5.5
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

#### **SEQUENCE**

TPEPCQIDDE DVRCVCNFTH PQPDWSSALQ CVAAVEVEIR GGGRSLEEFL LKSASANPKQ YADMLKALRL RRLTVGAARV PAQILALVLH ALGFSRLKEL TLEDLEVTGQ VPPPLQETPG PALTTLRLRN VSWATGGAWL GELQQWLQPS LKVLKVAQAS SLAFPCAQLR AFPALTTLDL SDNPGLGERG LTAALCPRKF PALEDLALRN AGMETPSGVC AALAGAGVQP HRLDLSHNSL RATAAGAREC VWPAALSSLN LSFAELEQVP KGLPPKLTVL DLSCNKLNRA PRPEELPAVD DLTLEGNPYM DPEALQHQED PMASGVVPPC ARSALTMGVS GTLALLQGAR GFA

#### USAGE


Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at - 80°C for 12 months.

#### STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

www.sunlongbiotech.com | sales@sunlongbiotech.com | +86 571 56623320