Recombinant Nitric Oxide Synthase 2, Inducible (NOS2) Instruction Manual

SIPA190Hu03

Homo sapiens (Human)

Source Prokaryotic expression

Host E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationCytoplasmPredicted Molecular Mass61.7kDa

Accurate Molecular Mass 62kDa(Analysis of differences refer to the manual)

Residues & Tags Met1~Glu504 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Traits Freeze-dried powder

Purity > 90% Isoelectric Point 7.0

Applications Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

MACPWKFLFK TKFHQYAMNG EKDINNNVEK APCATSSPVT QDDLQYHNLS KQQNESPQPL VETGKKSPES LVKLDATPLS SPRHVRIKNW GSGMTFQDTL HHKAKGILTC RSKSCLGSIM TPKSLTRGPR DKPTPPDELL PQAIEFVNQY YGSFKEAKIE EHLARVEAVT KEIETTGTYQ LTGDELIFAT KQAWRNAPRC IGRIQWSNLQ VFDARSCSTA REMFEHICRH VRYSTNNGNI RSAITVFPQR SDGKHDFRVW NAQLIRYAGY QMPDGSIRGD PANVEFTQLC IDLGWKPKYG RFDVVPLVLQ ANGRDPELFE IPPDLVLEVA MEHPKYEWFR ELELKWYALP AVANMLLEVG GLEFPGCPFN GWYMGTEIGV RDFCDVQRYN ILEEVGRRMG LETHKLASLW KDQAVVEINI AVLHSFQKQN VTIMDHHSAA ESFMKYMQNE YRSRGGCPAD WIWLVPPMSG SITPVFHQEM LNYVLSPFYY YQVEAWKTHV WQDE

USAGE

Reconstitute in ddH2O to a concentration of 0.1-1.0 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.





[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.