

Recombinant Phospholipase A2 Group VII (LpPLA2) Instruction Manual

SIPA201Hu01

Homo sapiens (Human)

Source	Prokaryotic expression
Host	E.coli
Endotoxin Level	<1.0EU per 1µg (determined by the LAL method)
Subcellular Location	Extracellular matrix
Predicted Molecular Mass	50.0kDa
Accurate Molecular Mass	50kDa(Analysis of differences refer to the manual)
Residues & Tags	Phe22~Asn441 with N-terminal His Tag
Buffer Formulation	20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose.
Traits	Freeze-dried powder
Purity	> 95%
Isoelectric Point	7.1
Applications	Positive Control; Immunogen; SDS-PAGE; WB.

SEQUENCE

```
FDWQYINPV AHMKSSAWVN KIQVLMMAAS
FGQTKIPRGN GPYSVGCTDL MFDHTNKGTF LRLYYPSQDN DRLDTLWIPN
KEYFWGLSKF LGTHWLMGNI LRLFLGSMIT PANWNSPLRP GEKYPLVVFS
HGLGAFRTLY SAIGIDLASH GFIVAAVEHR DRASATYYF KDQSAAEIGD
KSWLYLRTLK QEEETHIRNE QVRQRAKECS QALSLILDID HGKPVKNALD
LKFDMEQLKD SIDREKIAVI GHSFGGATVI QTLSEDQRFR CGIALDAWMF
PLGDEVYSRI PQPLFFINSE YFQYPANIIK MKKCYSPPKE RKMITIRGSV
HQNFAFDFTA TGKIIGHMLK LKGDIDSIVA IDLSNKASLA FLQKHLGLHK
DFDQWDCLIE GDDENLIPGT NINTTNQHIM LQNSSGIEKY N
```

USAGE

Reconstitute in ddH₂O to a concentration of 0.1-0.15 mg/mL. Do not vortex.


STORAGE

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

STABILITY

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Image



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.

