# **Recombinant Ceruloplasmin (CP) Instruction Manual**

# SIPA316Mu01

# Mus musculus (Mouse)

**Source** Prokaryotic expression

**Host** E.coli

Endotoxin Level <1.0EU per 1µg (determined by the LAL method)

Subcellular LocationSecretedPredicted Molecular Mass42.1kDa

Accurate Molecular Mass 42kDa(Analysis of differences refer to the manual)

**Residues & Tags** Tyr729~Gly1061 with N-terminal His Tag

Buffer Formulation 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

**Traits** Freeze-dried powder

Purity > 97% Isoelectric Point 6.5

**Applications** Positive Control; Immunogen; SDS-PAGE; WB.

# **SEQUENCE**

YY VAAVEVEWDY SPSRAWEKEL
HHLQEQNVSN VFLDKEEFFI GSKYKKVVYR QFTDSSFREQ VKRRAEDEHL
GILGPPIHAN VGDKVKVVFK NMATRPYSIH AHGVKTESST VVPTLPGEVR
TYTWQIPERS GAGREDSACI PWAYYSTVDR VKDLYSGLIG PLIVCRKSYV
KVFSPKKKME FFLLFLVFDE NESWYLDDNI KTYSEHPEKV NKDNEEFLES
NKMHAINGKM FGNLQGLTMH VKDEVNWYVM GMGNEIDLHT VHFHGHSFQY
KHRGVYSSDV FDLFPGTYQT LEMFPQTPGT WLLHCHVTDH VHAGMATTYT
VLPVEQETKS G

#### **USAGE**

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


#### **STORAGE**

Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

#### **STABILITY**

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.





SDS-PAGE Image

### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.